

ERA - EDTA Registry

Survival data for Austria
2002



Table of Contents

A5 SURVIVAL PROBABILITY (residents only)

Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.1	4
90-day, one-, two- and five-year survival probabilities (cohort 1993-1997): Incident RRT patients, unadjusted <i>from day 1 to 90 days, one, two and five years, by age, gender and primary diagnosis</i>	
Table A.5.2	5
90-day, one-, two- and five-year survival probabilities (cohort 1993-1997): Incident RRT patients, adjusted <i>from day 1 to 90 days, one, two and five years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.3	6
One-, two- and five-year survival probabilities (cohort 1993-1997): Incident RRT patients, unadjusted <i>from day 91 to one, two and five years + 90 days, by age, gender and primary diagnosis</i>	
Table A.5.4	7
One-, two- and five-year survival probabilities (cohort 1993-1997): Incident RRT patients, adjusted <i>from day 91 to one, two and five years + 90 days, adjusted for age, gender and primary diagnosis</i>	
Table A.5.5	8
90-day, one- and two-year survival probabilities (cohort 1996-2000): Incident RRT patients, unadjusted <i>from day 1 to 90 days, one and two years, by age, gender and primary diagnosis</i>	
Table A.5.6	9
90-day, one- and two-year survival probabilities (cohort 1996-2000): Incident RRT patients, adjusted <i>from day 1 to 90 days, one and two years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.7	10
One- and two-year survival probabilities (cohort 1996-2000): Incident RRT patients, unadjusted <i>from day 91 to one and two years + 90 days, by age, gender and primary diagnosis</i>	
Table A.5.8	11
One- and two-year survival probabilities (cohort 1996-2000): Incident RRT patients, adjusted <i>from day 91 to one and two years + 90 days, adjusted for age, gender and primary diagnosis</i>	

Patient Survival on Dialysis

Table A.5.9	12
90-day, one-, two- and five-year survival probabilities (cohort 1993-1997): Incident dialysis patients, unadjusted <i>from day 1 to 90 days, one, two and five years, by age, gender and primary diagnosis</i>	
Table A.5.10	13
90-day, one-, two- and five-year survival probabilities (cohort 1993-1997): Incident dialysis patients, adjusted <i>from day 1 to 90 days, one, two and five years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.11	14
One-, two- and five-year survival probabilities (cohort 1993-1997): Incident dialysis patients, unadjusted <i>from day 91 to one, two and five years + 90 days, by age, gender and primary diagnosis</i>	
Table A.5.12	15
One-, two- and five-year survival probabilities (cohort 1993-1997): Incident dialysis patients, adjusted <i>from day 91 to one, two and five years + 90 days, adjusted for age, gender and primary diagnosis</i>	
Table A.5.13	16
90-day, one- and two-year survival probabilities (cohort 1996-2000): Incident dialysis patients, unadjusted <i>from day 1 to 90 days, one and two years, by age, gender and primary diagnosis</i>	
Table A.5.14	17
90-day, one- and two-year survival probabilities (cohort 1996-2000): Incident dialysis patients, adjusted <i>from day 1 to 90 days, one and two years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.15	18
One- and two-year survival probabilities (cohort 1996-2000): Incident dialysis patients, unadjusted <i>from day 91 to one and two years + 90 days, by age, gender and primary diagnosis</i>	
Table A.5.16	19
One- and two-year survival probabilities (cohort 1996-2000): Incident dialysis patients, adjusted <i>from day 91 to one and two years + 90 days, adjusted for age, gender and primary diagnosis</i>	

Patient Survival after First Transplant (cadaver donor)

Table A.5.17 20
One-, two- and five-year survival probabilities (cohort 1993-1997): First transplant patients (cadaver donor), unadjusted from day of transplant to one, two and five years, by age, gender and primary diagnosis

Table A.5.18 21
One-, two- and five-year survival probabilities (cohort 1993-1997): First transplant patients (cadaver donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

Table A.5.19 22
One- and two-year survival probabilities (cohort 1996-2000): First transplant patients (cadaver donor), unadjusted from day of transplant to one and two years, by age, gender and primary diagnosis

Table A.5.20 23
One- and two-year survival probabilities (cohort 1996-2000): First transplant patients (cadaver donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

Patient Survival after First Transplant (living donor)

Table A.5.21 24
One-, two- and five-year survival probabilities (cohort 1993-1997): First transplant patients (living donor), unadjusted from day of transplant to one, two and five years, by age, gender and primary diagnosis

Table A.5.22 25
One-, two- and five-year survival probabilities (cohort 1993-1997): First transplant patients (living donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

Table A.5.23 26
One- and two-year survival probabilities (cohort 1996-2000): First transplant patients (living donor), unadjusted from day of transplant to one and two years, by age, gender and primary diagnosis

Table A.5.24 27
One- and two-year survival probabilities (cohort 1996-2000): First transplant patients (living donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

Graft Survival after First Transplant (cadaver donor)

Table A.5.25 28
One-, two- and five-year survival probabilities (cohort 1993-1997): First graft (cadaver donor), unadjusted from day of transplant to one, two and five years, by age, gender and primary diagnosis

Table A.5.26 29
One-, two- and five-year survival probabilities (cohort 1993-1997): First graft (cadaver donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

Table A.5.27 30
One- and two-year survival probabilities (cohort 1996-2000): First graft (cadaver donor), unadjusted from day of transplant to one and two years, by age, gender and primary diagnosis

Table A.5.28 31
One- and two-year survival probabilities (cohort 1996-2000): First graft (cadaver donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

Graft Survival after First Transplant (living donor)

Table A.5.29 32
One-, two- and five-year survival probabilities (cohort 1993-1997): First graft (living donor), unadjusted from day of transplant to one, two and five years, by age, gender and primary diagnosis

Table A.5.30 33
One-, two- and five-year survival probabilities (cohort 1993-1997): First graft (living donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

Table A.5.31 34
One- and two-year survival probabilities (cohort 1996-2000): First graft (living donor), unadjusted from day of transplant to one and two years, by age, gender and primary diagnosis

Table A.5.32 35
One- and two-year survival probabilities (cohort 1996-2000): First graft (living donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.1

90-day, one-, two- and five-year survival probabilities (cohort 1993-1997): Incident RRT patients, unadjusted
from day 1 to 90 days, one, two and five years, by age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES				
0-19	99.2 (98.4 - 99.6)	96.6 (95.3 - 97.6)	94.6 (93.1 - 95.9)	92.1 (90.3 - 93.6)
20-44	98.7 (98.4 - 98.9)	95.4 (94.9 - 95.8)	91.5 (90.8 - 92.0)	83.1 (82.4 - 83.9)
45-64	95.9 (95.6 - 96.2)	87.3 (86.9 - 87.8)	77.6 (77.1 - 78.1)	56.3 (55.9 - 56.8)
65-74	91.6 (91.1 - 92.0)	75.8 (75.2 - 76.4)	59.8 (59.3 - 60.3)	27.9 (27.7 - 28.2)
75+	86.0 (85.2 - 86.7)	65.0 (64.2 - 65.7)	46.6 (46.1 - 47.2)	14.8 (14.7 - 15.0)
Male	93.7 (93.4 - 93.9)	82.1 (81.7 - 82.5)	70.3 (69.9 - 70.7)	46.7 (46.4 - 47.0)
Female	93.7 (93.4 - 94.0)	82.0 (81.5 - 82.5)	70.4 (69.9 - 70.8)	47.2 (46.8 - 47.5)
Diabetes	93.5 (93.0 - 94.0)	77.8 (77.1 - 78.5)	60.0 (59.3 - 60.6)	30.1 (29.8 - 30.4)
Hypertension / renal vascular disease	91.3 (90.7 - 91.9)	77.3 (76.5 - 78.1)	63.0 (62.2 - 63.7)	33.9 (33.5 - 34.3)
Glomerulonephritis	96.4 (96.0 - 96.8)	90.2 (89.6 - 90.8)	83.1 (82.4 - 83.8)	66.4 (65.6 - 67.1)
Other cause	93.5 (93.2 - 93.9)	82.5 (82.1 - 82.9)	72.4 (72.0 - 72.9)	51.0 (50.6 - 51.3)
All	93.7 (93.5 - 93.9)	82.1 (81.8 - 82.4)	70.3 (70.0 - 70.6)	46.9 (46.6 - 47.1)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA				
0-19	100	95.8 (88.0-98.6)	92.9 (84.8-96.8)	91.5 (83.1-95.8)
20-44	98.5 (97.3-99.1)	96.4 (94.8-97.5)	93.0 (91.0-94.5)	83.8 (81.3-85.9)
45-64	94.8 (93.7-95.7)	84.2 (82.7-85.6)	73.4 (71.8-74.9)	48.0 (46.8-49.1)
65-74	92.1 (90.6-93.4)	72.5 (70.6-74.2)	55.2 (53.6-56.7)	21.7 (21.2-22.2)
75+	87.2 (84.6-89.4)	62.3 (59.7-64.7)	39.0 (37.5-40.6)	11.0 (10.7-11.2)
Male	94.1 (93.2-94.9)	81.3 (80.0-82.5)	68.8 (67.5-70.0)	43.9 (43.0-44.7)
Female	93.1 (92.0-94.1)	78.5 (77.0-80.0)	65.1 (63.6-66.5)	40.1 (39.2-41.0)
Diabetes	92.5 (91.0-93.8)	73.4 (71.5-75.2)	55.0 (53.4-56.5)	23.7 (23.2-24.3)
Hypertension / renal vascular disease	93.6 (91.3-95.3)	77.5 (74.6-80.2)	62.8 (60.1-65.3)	32.4 (31.1-33.7)
Glomerulonephritis	96.6 (95.0-97.7)	90.6 (88.4-92.4)	84.3 (81.8-86.4)	68.8 (66.3-71.2)
Other cause	93.5 (92.4-94.4)	81.4 (79.9-82.7)	70.1 (68.6-71.5)	47.3 (46.2-48.3)
All	93.7 (93.0-94.4)	80.1 (79.2-81.1)	67.2 (66.3-68.2)	42.3 (41.7-42.9)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.2

90-day, one-, two- and five-year survival probabilities (cohort 1993-1997): Incident RRT patients, adjusted from day 1 to 90 days, one, two and five years, adjusted for age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES				
0-19	99.1 (98.5 - 99.8)	96.0 (94.7 - 97.4)	93.7 (92.1 - 95.5)	90.8 (88.8 - 92.8)
20-44	98.7 (98.4 - 98.9)	95.3 (94.8 - 95.8)	91.3 (90.6 - 91.9)	82.6 (81.7 - 83.5)
45-64	96.1 (95.8 - 96.4)	87.7 (87.2 - 88.2)	78.2 (77.5 - 78.8)	56.5 (55.7 - 57.3)
65-74	91.9 (91.5 - 92.4)	76.8 (76.1 - 77.6)	61.1 (60.2 - 62.0)	28.6 (27.8 - 29.4)
75+	86.1 (85.2 - 86.9)	65.3 (64.1 - 66.4)	46.9 (45.7 - 48.2)	14.5 (13.7 - 15.5)
Male	95.1 (94.9 - 95.4)	85.7 (85.3 - 86.1)	75.0 (74.5 - 75.6)	49.0 (48.3 - 49.7)
Female	95.4 (95.2 - 95.7)	86.4 (85.9 - 86.9)	76.3 (75.7 - 76.9)	51.5 (50.7 - 52.3)
Diabetes	94.7 (94.3 - 95.1)	81.1 (80.3 - 81.9)	64.2 (63.2 - 65.2)	31.0 (29.9 - 32.1)
Hypertension / renal vascular disease	94.9 (94.5 - 95.3)	85.9 (85.2 - 86.6)	75.6 (74.7 - 76.5)	48.3 (47.1 - 49.6)
Glomerulonephritis	96.5 (96.1 - 96.9)	90.1 (89.4 - 90.8)	82.3 (81.4 - 83.3)	61.8 (60.6 - 63.1)
Other cause	94.9 (94.7 - 95.2)	85.9 (85.4 - 86.3)	76.8 (76.2 - 77.4)	54.1 (53.3 - 54.8)
All	95.3 (95.1 - 95.4)	86.0 (85.6 - 86.3)	75.5 (75.1 - 76.0)	50.0 (49.5 - 50.6)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA				
0-19				
20-44	98.4 (97.5-99.3)	96.2 (94.8-97.7)	92.7 (90.8-94.7)	83.1 (80.3-86.0)
45-64	95.2 (94.2-96.2)	85.1 (83.5-86.8)	74.9 (73.0-77.0)	49.4 (47.1-51.9)
65-74	92.7 (91.4-94.1)	75.3 (73.0-77.6)	58.9 (56.3-61.6)	24.7 (22.4-27.3)
75+	88.3 (85.8-90.8)	65.0 (61.3-68.9)	42.1 (38.3-46.4)	12.8 (10.2-16.0)
Male	95.4 (94.7-96.1)	84.8 (83.5-86.1)	73.2 (71.6-74.9)	45.2 (43.2-47.3)
Female	95.6 (94.8-96.3)	85.3 (83.9-86.7)	74.1 (72.4-75.9)	46.3 (44.0-48.8)
Diabetes	94.3 (93.2-95.4)	79.0 (77.0-81.1)	62.1 (59.5-64.7)	27.2 (24.7-30.0)
Hypertension / renal vascular disease	96.1 (94.8-97.4)	85.6 (83.2-88.0)	74.4 (71.3-77.6)	44.6 (40.6-49.0)
Glomerulonephritis	96.0 (94.4-97.6)	88.5 (85.9-91.2)	80.7 (77.4-84.1)	59.5 (55.3-64.0)
Other cause	94.7 (93.8-95.6)	84.4 (82.9-85.9)	73.9 (72.0-75.8)	48.5 (46.2-50.9)
All	95.5 (94.9-96.0)	85.0 (84.0-85.9)	73.5 (72.3-74.8)	45.6 (44.1-47.2)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.3

One-, two- and five-year survival probabilities (cohort 1993-1997): Incident RRT patients, unadjusted from day 91 to one, two and five years + 90 days, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	96.4 (95.0 - 97.5)	94.8 (93.1 - 96.1)	92.1 (90.2 - 93.7)
20-44	95.5 (95.0 - 95.9)	91.6 (91.0 - 92.2)	83.0 (82.2 - 83.7)
45-64	88.2 (87.7 - 88.6)	78.6 (78.1 - 79.1)	56.6 (56.2 - 57.1)
65-74	78.3 (77.7 - 78.9)	61.3 (60.8 - 61.9)	28.4 (28.2 - 28.7)
75+	69.8 (69.0 - 70.7)	49.8 (49.1 - 50.4)	15.3 (15.1 - 15.4)
Male	84.1 (83.7 - 84.5)	72.0 (71.6 - 72.4)	47.5 (47.2 - 47.8)
Female	84.2 (83.7 - 84.6)	72.2 (71.7 - 72.7)	48.2 (47.8 - 48.6)
Diabetes	77.9 (77.1 - 78.6)	60.1 (59.5 - 60.8)	29.6 (29.3 - 29.9)
Hypertension / renal vascular disease	80.4 (79.6 - 81.2)	65.2 (64.4 - 66.0)	35.3 (34.9 - 35.8)
Glomerulonephritis	91.5 (90.8 - 92.1)	84.5 (83.7 - 85.2)	67.0 (66.2 - 67.8)
Other cause	85.2 (84.8 - 85.7)	74.7 (74.2 - 75.2)	52.3 (51.9 - 52.6)
All	84.1 (83.8 - 84.4)	72.1 (71.8 - 72.4)	47.8 (47.5 - 48.0)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19	92.3 (82.1-96.8)	90.4 (79.9-95.5)	88.4 (77.8-94.1)
20-44	96.6 (95.0-97.7)	93.6 (91.6-95.1)	83.5 (81.0-85.7)
45-64	85.9 (84.3-87.3)	74.7 (73.1-76.3)	48.9 (47.7-50.1)
65-74	73.5 (71.6-75.4)	55.5 (53.9-57.1)	21.6 (21.1-22.1)
75+	63.3 (60.5-65.9)	40.4 (38.7-42.2)	11.6 (11.3-11.9)
Male	82.6 (81.3-83.9)	70.2 (68.9-71.5)	44.5 (43.6-45.4)
Female	79.8 (78.2-81.3)	66.0 (64.5-67.5)	40.6 (39.6-41.6)
Diabetes	73.6 (71.6-75.5)	55.5 (53.9-57.2)	23.3 (22.7-23.9)
Hypertension / renal vascular disease	78.3 (75.3-81.1)	63.0 (60.2-65.7)	32.6 (31.2-34.0)
Glomerulonephritis	92.2 (90.1-93.9)	85.8 (83.3-88.0)	69.7 (67.1-72.1)
Other cause	83.3 (81.7-84.7)	71.7 (70.2-73.2)	48.4 (47.3-49.6)
All	81.4 (80.4-82.4)	68.4 (67.4-69.4)	42.9 (42.2-43.6)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.4

One-, two- and five-year survival probabilities (cohort 1993-1997): Incident RRT patients, adjusted from day 91 to one, two and five years + 90 days, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	95.7 (94.2 - 97.3)	93.8 (92.0 - 95.7)	90.7 (88.6 - 92.9)
20-44	95.4 (94.9 - 95.9)	91.5 (90.8 - 92.2)	82.7 (81.8 - 83.6)
45-64	88.6 (88.1 - 89.1)	79.2 (78.5 - 79.9)	57.3 (56.5 - 58.1)
65-74	79.2 (78.5 - 80.0)	62.6 (61.7 - 63.5)	29.4 (28.6 - 30.3)
75+	70.1 (68.8 - 71.3)	49.8 (48.5 - 51.2)	15.1 (14.1 - 16.1)
Male	87.1 (86.7 - 87.5)	76.1 (75.6 - 76.7)	49.6 (48.9 - 50.3)
Female	87.9 (87.4 - 88.3)	77.4 (76.8 - 78.1)	52.2 (51.4 - 53.1)
Diabetes	80.9 (80.0 - 81.7)	63.9 (62.9 - 65.0)	30.3 (29.2 - 31.4)
Hypertension / renal vascular disease	87.6 (87.0 - 88.3)	76.6 (75.7 - 77.5)	49.2 (48.0 - 50.5)
Glomerulonephritis	91.2 (90.5 - 91.9)	83.5 (82.6 - 84.4)	62.5 (61.2 - 63.8)
Other cause	87.9 (87.5 - 88.3)	78.4 (77.8 - 79.0)	55.1 (54.4 - 55.9)
All	87.4 (87.1 - 87.7)	76.7 (76.2 - 77.1)	50.7 (50.1 - 51.2)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	96.5 (95.1-97.9)	93.4 (91.5-95.3)	83.1 (80.3-86.1)
45-64	87.0 (85.4-88.6)	76.4 (74.4-78.5)	50.8 (48.4-53.3)
65-74	76.3 (74.0-78.7)	59.5 (56.8-62.3)	25.3 (22.8-28.0)
75+	66.0 (62.1-70.2)	43.6 (39.5-48.2)	13.7 (10.9-17.2)
Male	85.8 (84.5-87.1)	74.3 (72.6-76.0)	46.1 (44.0-48.3)
Female	86.1 (84.7-87.5)	74.5 (72.7-76.4)	46.8 (44.3-49.3)
Diabetes	78.8 (76.6-81.0)	62.0 (59.4-64.8)	26.5 (23.9-29.4)
Hypertension / renal vascular disease	85.9 (83.4-88.4)	74.1 (70.8-77.5)	44.8 (40.7-49.3)
Glomerulonephritis	90.6 (88.1-93.1)	82.5 (79.3-85.9)	60.7 (56.5-65.3)
Other cause	85.9 (84.4-87.4)	75.1 (73.2-77.0)	49.7 (47.4-52.2)
All	85.9 (84.9-86.9)	74.3 (73.1-75.6)	46.3 (44.7-48.0)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.5

90-day, one- and two-year survival probabilities (cohort 1996-2000): Incident RRT patients, unadjusted from day 1 to 90 days, one and two years, by age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES			
0-19	99.3 (98.5 - 99.6)	96.8 (95.5 - 97.7)	95.8 (94.4 - 96.9)
20-44	98.5 (98.3 - 98.8)	95.1 (94.7 - 95.5)	91.6 (91.1 - 92.2)
45-64	96.0 (95.7 - 96.3)	87.5 (87.1 - 87.9)	78.4 (77.9 - 78.9)
65-74	90.9 (90.5 - 91.3)	75.7 (75.2 - 76.2)	60.8 (60.3 - 61.2)
75+	86.7 (86.1 - 87.2)	66.4 (65.8 - 67.0)	48.4 (47.9 - 48.8)
Male	93.3 (93.0 - 93.5)	81.5 (81.2 - 81.8)	69.8 (69.4 - 70.1)
Female	92.9 (92.6 - 93.2)	80.8 (80.3 - 81.2)	69.3 (68.9 - 69.7)
Diabetes	93.5 (93.1 - 93.9)	79.1 (78.5 - 79.6)	62.5 (61.9 - 63.1)
Hypertension / renal vascular disease	91.3 (90.8 - 91.9)	76.6 (75.9 - 77.2)	62.7 (62.1 - 63.4)
Glomerulonephritis	96.3 (95.9 - 96.7)	90.5 (89.9 - 91.0)	83.7 (83.0 - 84.3)
Other cause	92.6 (92.3 - 92.8)	80.9 (80.5 - 81.3)	70.6 (70.3 - 71.0)
All	93.1 (92.9 - 93.3)	81.2 (80.9 - 81.5)	69.6 (69.3 - 69.9)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA			
0-19	100	96.9 (88.7-99.2)	95.4 (87.0-98.4)
20-44	97.6 (96.2-98.5)	94.8 (93.1-96.2)	92.2 (90.2-93.8)
45-64	94.8 (93.7-95.7)	83.3 (81.8-84.7)	72.5 (70.9-73.9)
65-74	89.7 (88.2-91.1)	72.4 (70.7-74.1)	55.5 (54.1-57.0)
75+	85.3 (83.2-87.2)	62.8 (60.6-64.8)	43.7 (42.2-45.2)
Male	93.3 (92.4-94.1)	80.2 (79.0-81.3)	68.1 (66.9-69.3)
Female	90.5 (89.3-91.6)	76.2 (74.7-77.6)	62.8 (61.4-64.1)
Diabetes	90.8 (89.3-92.0)	73.5 (71.8-75.1)	55.3 (53.9-56.7)
Hypertension / renal vascular disease	93.1 (91.2-94.6)	79.4 (77.0-81.7)	62.2 (60.0-64.4)
Glomerulonephritis	96.2 (94.6-97.4)	89.2 (86.9-91.1)	83.7 (81.2-85.9)
Other cause	91.5 (90.3-92.5)	78.3 (76.8-79.7)	69.1 (67.7-70.5)
All	92.2 (91.4-92.8)	78.6 (77.6-79.5)	66.0 (65.1-66.8)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.6

90-day, one- and two-year survival probabilities (cohort 1996-2000): Incident RRT patients, adjusted from day 1 to 90 days, one and two years, adjusted for age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES			
0-19	99.2 (98.6 - 99.8)	96.4 (95.2 - 97.7)	95.4 (94.0 - 96.8)
20-44	98.5 (98.2 - 98.8)	95.0 (94.5 - 95.5)	91.5 (90.8 - 92.1)
45-64	96.1 (95.8 - 96.4)	87.9 (87.5 - 88.4)	79.0 (78.4 - 79.6)
65-74	91.3 (90.9 - 91.7)	76.8 (76.2 - 77.5)	62.3 (61.5 - 63.0)
75+	86.8 (86.2 - 87.5)	66.9 (66.0 - 67.8)	49.0 (48.0 - 49.9)
Male	95.0 (94.8 - 95.2)	85.9 (85.5 - 86.2)	75.8 (75.3 - 76.3)
Female	95.1 (94.8 - 95.3)	86.0 (85.6 - 86.5)	76.6 (76.0 - 77.1)
Diabetes	94.8 (94.5 - 95.2)	82.8 (82.1 - 83.4)	67.8 (66.9 - 68.6)
Hypertension / renal vascular disease	95.0 (94.7 - 95.4)	85.8 (85.2 - 86.4)	76.0 (75.3 - 76.8)
Glomerulonephritis	96.5 (96.1 - 96.9)	90.7 (90.1 - 91.4)	83.7 (82.8 - 84.5)
Other cause	94.4 (94.2 - 94.7)	85.2 (84.8 - 85.6)	76.3 (75.8 - 76.8)
All	95.0 (94.9 - 95.2)	85.9 (85.7 - 86.2)	76.1 (75.7 - 76.5)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	97.5 (96.3-98.6)	94.6 (93.0-96.3)	91.9 (89.9-93.9)
45-64	95.1 (94.2-96.1)	84.4 (82.8-86.0)	74.0 (72.0-76.0)
65-74	90.8 (89.4-92.2)	75.3 (73.2-77.5)	59.4 (57.0-61.9)
75+	86.2 (84.0-88.5)	65.3 (62.2-68.5)	46.8 (43.5-50.3)
Male	94.9 (94.2-95.6)	84.3 (83.1-85.5)	73.3 (71.8-74.8)
Female	94.3 (93.5-95.1)	84.7 (83.4-86.0)	74.4 (72.7-76.0)
Diabetes	93.3 (92.2-94.4)	79.8 (78.0-81.6)	63.8 (61.6-66.1)
Hypertension / renal vascular disease	95.8 (94.7-96.9)	87.2 (85.3-89.1)	74.5 (71.9-77.2)
Glomerulonephritis	95.7 (94.1-97.3)	87.3 (84.6-90.0)	80.5 (77.3-83.8)
Other cause	93.2 (92.3-94.2)	82.2 (80.7-83.8)	73.8 (72.0-75.6)
All	94.6 (94.0-95.1)	84.4 (83.5-85.3)	73.7 (72.6-74.8)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.7

One- and two-year survival probabilities (cohort 1996-2000): Incident RRT patients, unadjusted
from day 91 to one and two years + 90 days, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	97.2 (95.9 - 98.1)	96.2 (94.7 - 97.3)
20-44	95.6 (95.1 - 96.0)	91.9 (91.3 - 92.4)
45-64	88.5 (88.1 - 88.9)	79.4 (78.9 - 79.9)
65-74	79.2 (78.6 - 79.7)	63.0 (62.5 - 63.5)
75+	71.1 (70.4 - 71.8)	51.3 (50.8 - 51.8)
Male	84.0 (83.7 - 84.3)	71.7 (71.3 - 72.0)
Female	83.6 (83.1 - 84.0)	71.7 (71.2 - 72.1)
Diabetes	79.7 (79.1 - 80.3)	62.8 (62.2 - 63.4)
Hypertension / renal vascular disease	79.9 (79.2 - 80.6)	64.9 (64.2 - 65.6)
Glomerulonephritis	92.1 (91.5 - 92.6)	85.0 (84.3 - 85.7)
Other cause	84.3 (84.0 - 84.7)	73.6 (73.1 - 74.0)
All	83.8 (83.6 - 84.1)	71.7 (71.4 - 72.0)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19	95.4 (83.8-98.8)	95.4 (83.8-98.8)
20-44	96.3 (94.6-97.4)	93.1 (91.0-94.7)
45-64	84.7 (83.1-86.1)	73.3 (71.7-74.9)
65-74	75.9 (74.0-77.7)	57.4 (55.8-59.0)
75+	67.4 (65.0-69.7)	44.4 (42.7-46.0)
Male	82.1 (80.8-83.3)	68.7 (67.5-70.0)
Female	80.1 (78.5-81.5)	65.4 (63.9-66.9)
Diabetes	75.5 (73.7-77.2)	56.6 (55.0-58.1)
Hypertension / renal vascular disease	81.0 (78.4-83.3)	62.5 (60.1-64.8)
Glomerulonephritis	91.1 (88.9-92.9)	84.4 (81.8-86.7)
Other cause	82.2 (80.6-83.6)	71.4 (69.8-72.9)
All	81.3 (80.3-82.2)	67.4 (66.4-68.3)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.8

One- and two-year survival probabilities (cohort 1996-2000): Incident RRT patients, adjusted from day 91 to one and two years + 90 days, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	96.8 (95.5 - 98.0)	95.7 (94.2 - 97.2)
20-44	95.5 (95.1 - 96.0)	91.8 (91.2 - 92.4)
45-64	88.9 (88.5 - 89.4)	80.2 (79.6 - 80.8)
65-74	80.2 (79.6 - 80.9)	64.8 (64.1 - 65.6)
75+	71.5 (70.6 - 72.4)	52.2 (51.2 - 53.3)
Male	87.6 (87.3 - 88.0)	77.2 (76.7 - 77.7)
Female	87.9 (87.5 - 88.3)	78.3 (77.7 - 78.8)
Diabetes	83.0 (82.4 - 83.7)	68.0 (67.1 - 68.8)
Hypertension / renal vascular disease	87.7 (87.1 - 88.2)	77.3 (76.5 - 78.1)
Glomerulonephritis	92.1 (91.5 - 92.7)	84.9 (84.1 - 85.7)
Other cause	87.7 (87.3 - 88.0)	78.4 (77.9 - 78.9)
All	87.8 (87.5 - 88.0)	77.6 (77.3 - 78.0)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	96.2 (94.7-97.6)	92.9 (90.9-94.9)
45-64	85.9 (84.3-87.5)	75.4 (73.4-77.4)
65-74	78.7 (76.6-80.9)	62.0 (59.4-64.6)
75+	70.0 (66.8-73.4)	48.3 (44.8-52.1)
Male	85.8 (84.6-87.0)	74.1 (72.6-75.7)
Female	87.3 (86.1-88.6)	76.3 (74.6-78.1)
Diabetes	81.0 (79.1-82.9)	64.9 (62.6-67.3)
Hypertension / renal vascular disease	88.0 (86.0-90.0)	74.6 (71.8-77.4)
Glomerulonephritis	89.5 (87.0-92.2)	81.4 (78.1-84.8)
Other cause	85.3 (83.8-86.8)	75.5 (73.6-77.5)
All	86.4 (85.6-87.3)	75.0 (73.9-76.2)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Dialysis

Table A.5.9

90-day, one-, two- and five-year survival probabilities (cohort 1993-1997): Incident dialysis patients, unadjusted from day 1 to 90 days, one, two and five years, by age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES				
0-19	99.1 (98.1 - 99.6)	96.6 (95.0 - 97.7)	93.8 (91.5 - 95.5)	87.7 (84.0 - 90.6)
20-44	98.7 (98.4 - 98.9)	95.0 (94.4 - 95.5)	89.8 (89.0 - 90.5)	74.3 (73.1 - 75.4)
45-64	95.9 (95.6 - 96.2)	87.0 (86.5 - 87.5)	76.0 (75.4 - 76.5)	48.2 (47.7 - 48.7)
65-74	91.6 (91.2 - 92.1)	75.8 (75.3 - 76.4)	59.6 (59.1 - 60.1)	26.8 (26.5 - 27.0)
75+	86.0 (85.2 - 86.7)	65.1 (64.3 - 65.8)	46.8 (46.2 - 47.4)	14.8 (14.7 - 15.0)
Male	93.6 (93.3 - 93.8)	81.4 (81.0 - 81.8)	67.9 (67.4 - 68.3)	36.8 (36.5 - 37.0)
Female	93.6 (93.2 - 93.9)	81.4 (80.9 - 81.9)	68.3 (67.8 - 68.8)	38.5 (38.2 - 38.9)
Diabetes	93.5 (92.9 - 93.9)	77.3 (76.6 - 78.1)	58.0 (57.3 - 58.6)	23.2 (22.9 - 23.4)
Hypertension / renal vascular disease	91.3 (90.7 - 91.9)	77.2 (76.3 - 77.9)	62.1 (61.3 - 62.8)	29.4 (29.0 - 29.7)
Glomerulonephritis	96.4 (95.9 - 96.8)	89.6 (88.9 - 90.2)	80.6 (79.8 - 81.4)	54.4 (53.6 - 55.2)
Other cause	93.4 (93.1 - 93.7)	81.8 (81.3 - 82.2)	70.3 (69.8 - 70.8)	41.9 (41.5 - 42.2)
All	93.6 (93.3 - 93.8)	81.4 (81.1 - 81.7)	68.0 (67.7 - 68.4)	37.5 (37.3 - 37.7)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA				
0-19	100	100	92.3 (75.0-97.8)	84.9 (63.4-94.3)
20-44	98.6 (97.4-99.2)	96.2 (94.5-97.4)	91.7 (89.3-93.6)	75.5 (71.8-78.8)
45-64	94.9 (93.8-95.8)	84.0 (82.5-85.5)	72.4 (70.7-74.0)	40.4 (39.3-41.6)
65-74	92.2 (90.7-93.4)	72.8 (71.0-74.6)	55.5 (53.9-57.1)	20.7 (20.2-21.2)
75+	87.4 (84.8-89.5)	62.8 (60.3-65.3)	39.6 (38.0-41.3)	11.2 (11.0-11.5)
Male	94.2 (93.3-95.0)	81.0 (79.7-82.3)	67.0 (65.7-68.3)	34.2 (33.4-34.9)
Female	93.2 (92.0-94.2)	78.3 (76.7-79.8)	63.5 (62.0-65.0)	32.0 (31.2-32.8)
Diabetes	92.7 (91.2-93.9)	73.3 (71.4-75.1)	53.6 (52.0-55.2)	18.3 (17.8-18.7)
Hypertension / renal vascular disease	93.9 (91.6-95.5)	77.5 (74.5-80.1)	61.6 (58.9-64.3)	27.7 (26.5-28.9)
Glomerulonephritis	96.5 (94.9-97.6)	90.0 (87.7-91.9)	83.5 (80.7-85.8)	58.1 (55.2-60.8)
Other cause	93.5 (92.4-94.4)	81.3 (79.8-82.7)	68.6 (67.0-70.0)	38.4 (37.4-39.4)
All	93.8 (93.1-94.4)	79.9 (78.9-80.8)	65.5 (64.5-66.5)	33.2 (32.7-33.8)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Dialysis

Table A.5.10

90-day, one-, two- and five-year survival probabilities (cohort 1993-1997): Incident dialysis patients, adjusted from day 1 to 90 days, one, two and five years, adjusted for age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES				
0-19	98.9 (98.2 - 99.7)	96.1 (94.4 - 97.7)	92.9 (90.5 - 95.4)	85.2 (80.6 - 90.0)
20-44	98.6 (98.4 - 98.9)	94.9 (94.3 - 95.4)	89.5 (88.7 - 90.4)	71.8 (70.1 - 73.6)
45-64	96.0 (95.7 - 96.3)	87.4 (86.9 - 88.0)	76.7 (76.0 - 77.4)	47.6 (46.7 - 48.6)
65-74	91.9 (91.4 - 92.3)	76.7 (75.9 - 77.4)	60.7 (59.8 - 61.6)	27.0 (26.2 - 27.9)
75+	86.0 (85.2 - 86.9)	65.1 (64.0 - 66.3)	46.8 (45.6 - 48.1)	14.4 (13.5 - 15.3)
Male	94.6 (94.3 - 94.9)	84.2 (83.8 - 84.7)	72.3 (71.7 - 72.9)	41.6 (40.8 - 42.4)
Female	94.9 (94.6 - 95.2)	84.9 (84.4 - 85.5)	73.7 (73.0 - 74.4)	44.8 (43.9 - 45.8)
Diabetes	94.3 (93.8 - 94.7)	79.7 (78.8 - 80.5)	61.6 (60.5 - 62.7)	25.8 (24.7 - 26.9)
Hypertension / renal vascular disease	94.2 (93.7 - 94.6)	84.1 (83.3 - 84.8)	72.4 (71.4 - 73.5)	41.7 (40.4 - 43.0)
Glomerulonephritis	96.3 (95.8 - 96.7)	89.4 (88.6 - 90.2)	80.5 (79.5 - 81.6)	53.3 (51.8 - 54.9)
Other cause	94.4 (94.1 - 94.7)	84.4 (83.9 - 84.9)	74.3 (73.6 - 74.9)	47.0 (46.2 - 47.9)
All	94.7 (94.5 - 94.9)	84.5 (84.2 - 84.9)	72.9 (72.4 - 73.3)	42.9 (42.3 - 43.5)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA				
0-19				
20-44	98.5 (97.6-99.4)	96.1 (94.6-97.6)	91.5 (89.1-93.9)	72.1 (66.5-78.1)
45-64	95.2 (94.3-96.2)	84.9 (83.3-86.6)	73.9 (71.8-76.1)	40.8 (38.0-43.8)
65-74	92.6 (91.3-94.0)	75.1 (72.8-77.5)	58.5 (55.8-61.3)	22.3 (19.9-25.0)
75+	88.2 (85.7-90.7)	64.9 (61.1-68.8)	42.0 (38.1-46.3)	12.6 (10.1-15.8)
Male	95.0 (94.2-95.8)	83.6 (82.2-85.0)	70.7 (68.9-72.5)	37.0 (34.8-39.5)
Female	95.0 (94.2-95.9)	83.6 (82.1-85.1)	71.2 (69.2-73.2)	39.6 (37.0-42.3)
Diabetes	93.8 (92.6-95.1)	77.3 (75.1-79.5)	59.0 (56.4-61.9)	21.8 (19.2-24.7)
Hypertension / renal vascular disease	95.7 (94.3-97.2)	83.7 (81.0-86.5)	71.0 (67.5-74.6)	37.6 (33.2-42.5)
Glomerulonephritis	95.9 (94.2-97.5)	88.2 (85.5-91.0)	80.6 (77.2-84.2)	51.4 (46.2-57.2)
Other cause	94.2 (93.2-95.2)	83.3 (81.7-84.9)	71.5 (69.5-73.6)	41.1 (38.5-43.9)
All	95.0 (94.4-95.6)	83.5 (82.5-84.5)	70.8 (69.5-72.2)	38.0 (36.3-39.9)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Dialysis

Table A.5.11

One-, two- and five-year survival probabilities (cohort 1993-1997): Incident dialysis patients, unadjusted from day 91 to one, two and five years + 90 days, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	96.5 (94.8 - 97.7)	94.2 (91.8 - 95.9)	88.5 (84.6 - 91.5)
20-44	95.0 (94.4 - 95.5)	89.8 (89.0 - 90.5)	73.4 (72.2 - 74.6)
45-64	87.7 (87.2 - 88.2)	76.7 (76.1 - 77.2)	47.7 (47.2 - 48.2)
65-74	78.2 (77.6 - 78.8)	61.1 (60.5 - 61.6)	27.1 (26.8 - 27.3)
75+	70.0 (69.1 - 70.8)	49.9 (49.2 - 50.6)	15.3 (15.1 - 15.4)
Male	83.3 (82.9 - 83.7)	69.2 (68.7 - 69.6)	36.8 (36.5 - 37.1)
Female	83.4 (82.9 - 83.9)	69.7 (69.2 - 70.3)	38.8 (38.5 - 39.2)
Diabetes	77.3 (76.5 - 78.0)	57.8 (57.1 - 58.4)	22.2 (21.9 - 22.4)
Hypertension / renal vascular disease	80.1 (79.2 - 80.9)	64.0 (63.2 - 64.8)	30.1 (29.7 - 30.5)
Glomerulonephritis	90.7 (90.0 - 91.3)	81.6 (80.7 - 82.4)	54.1 (53.2 - 54.9)
Other cause	84.5 (84.0 - 84.9)	72.3 (71.8 - 72.8)	42.4 (42.1 - 42.8)
All	83.3 (83.0 - 83.7)	69.4 (69.1 - 69.8)	37.6 (37.4 - 37.8)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19	97.5 (84.1-99.6)	93.0 (75.9-98.1)	85.3 (62.8-94.7)
20-44	96.2 (94.4-97.4)	92.2 (89.8-94.1)	74.5 (70.5-78.0)
45-64	85.5 (83.9-86.9)	73.3 (71.6-75.0)	40.7 (39.5-41.9)
65-74	73.8 (71.8-75.6)	55.7 (54.0-57.3)	20.3 (19.8-20.8)
75+	63.9 (61.1-66.6)	41.0 (39.2-42.8)	11.9 (11.6-12.3)
Male	82.1 (80.7-83.3)	68.1 (66.7-69.4)	34.3 (33.5-35.1)
Female	79.6 (78.0-81.1)	64.3 (62.7-65.8)	32.3 (31.5-33.2)
Diabetes	73.2 (71.2-75.1)	53.7 (52.0-55.3)	17.1 (16.7-17.6)
Hypertension / renal vascular disease	78.0 (74.9-80.7)	61.7 (58.9-64.4)	27.6 (26.4-28.9)
Glomerulonephritis	91.8 (89.4-93.6)	84.6 (81.8-87.0)	58.2 (55.1-61.1)
Other cause	83.1 (81.6-84.6)	70.1 (68.5-71.7)	39.6 (38.5-40.7)
All	81.0 (80.0-82.0)	66.5 (65.4-67.5)	33.4 (32.9-34.0)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Dialysis

Table A.5.12

One-, two- and five-year survival probabilities (cohort 1993-1997): Incident dialysis patients, adjusted from day 91 to one, two and five years + 90 days, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	95.7 (93.9 - 97.6)	93.0 (90.4 - 95.6)	85.9 (81.3 - 90.8)
20-44	94.9 (94.3 - 95.5)	89.5 (88.6 - 90.4)	71.0 (69.2 - 72.9)
45-64	88.2 (87.6 - 88.7)	77.4 (76.7 - 78.1)	47.7 (46.7 - 48.7)
65-74	79.0 (78.2 - 79.7)	62.1 (61.2 - 63.0)	27.6 (26.8 - 28.5)
75+	70.0 (68.8 - 71.2)	49.7 (48.4 - 51.1)	14.9 (14.0 - 15.9)
Male	85.7 (85.2 - 86.1)	73.2 (72.6 - 73.8)	41.6 (40.8 - 42.4)
Female	86.4 (85.9 - 86.9)	74.6 (73.9 - 75.3)	45.0 (44.0 - 45.9)
Diabetes	79.3 (78.4 - 80.2)	61.0 (59.9 - 62.2)	24.7 (23.6 - 25.9)
Hypertension / renal vascular disease	85.8 (85.0 - 86.6)	73.3 (72.2 - 74.3)	42.0 (40.6 - 43.4)
Glomerulonephritis	90.5 (89.7 - 91.2)	81.4 (80.4 - 82.5)	53.2 (51.6 - 54.9)
Other cause	86.6 (86.1 - 87.0)	75.8 (75.1 - 76.4)	47.4 (46.5 - 48.3)
All	86.0 (85.6 - 86.3)	73.8 (73.3 - 74.3)	42.9 (42.3 - 43.6)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	96.0 (94.4-97.6)	91.8 (89.4-94.3)	69.8 (63.7-76.5)
45-64	86.6 (85.0-88.3)	75.1 (72.9-77.3)	41.6 (38.7-44.8)
65-74	76.1 (73.7-78.5)	58.9 (56.1-61.8)	22.3 (19.8-25.2)
75+	66.0 (62.0-70.2)	43.4 (39.3-48.0)	13.5 (10.7-17.0)
Male	84.4 (83.0-85.9)	71.6 (69.8-73.5)	37.4 (34.9-39.9)
Female	84.4 (82.8-86.0)	71.4 (69.3-73.5)	39.3 (36.5-42.2)
Diabetes	76.8 (74.5-79.3)	58.8 (55.9-61.7)	20.5 (17.9-23.5)
Hypertension / renal vascular disease	83.7 (80.9-86.7)	70.4 (66.7-74.3)	37.3 (32.7-42.4)
Glomerulonephritis	90.3 (87.7-92.9)	82.0 (78.5-85.7)	52.1 (46.6-58.2)
Other cause	84.8 (83.1-86.4)	72.6 (70.5-74.8)	41.7 (39.0-44.7)
All	84.3 (83.3-85.4)	71.4 (70.0-72.9)	38.1 (36.2-40.0)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Dialysis

Table A.5.13

90-day, one- and two-year survival probabilities (cohort 1996-2000): Incident dialysis patients, unadjusted from day 1 to 90 days, one and two years, by age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES			
0-19	99.1 (98.1 - 99.6)	96.4 (94.7 - 97.6)	95.1 (93.1 - 96.6)
20-44	98.6 (98.3 - 98.8)	94.7 (94.1 - 95.1)	90.1 (89.4 - 90.7)
45-64	95.9 (95.7 - 96.2)	87.3 (86.8 - 87.7)	77.0 (76.5 - 77.5)
65-74	91.1 (90.7 - 91.5)	76.0 (75.5 - 76.5)	60.9 (60.4 - 61.3)
75+	87.0 (86.4 - 87.6)	67.1 (66.5 - 67.7)	49.0 (48.6 - 49.5)
Male	93.3 (93.0 - 93.5)	81.0 (80.7 - 81.4)	67.9 (67.5 - 68.2)
Female	92.9 (92.6 - 93.2)	80.5 (80.1 - 80.9)	67.7 (67.3 - 68.2)
Diabetes	93.5 (93.1 - 93.9)	78.8 (78.2 - 79.4)	61.1 (60.6 - 61.7)
Hypertension / renal vascular disease	91.5 (91.0 - 92.1)	76.8 (76.1 - 77.4)	62.3 (61.7 - 62.9)
Glomerulonephritis	96.3 (95.9 - 96.7)	90.1 (89.4 - 90.7)	81.8 (81.0 - 82.5)
Other cause	92.5 (92.2 - 92.8)	80.4 (80.0 - 80.8)	68.8 (68.4 - 69.2)
All	93.1 (92.9 - 93.3)	80.8 (80.6 - 81.1)	67.8 (67.6 - 68.1)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA			
0-19	100	100	100
20-44	97.9 (96.5-98.7)	94.7 (92.8-96.1)	91.7 (89.4-93.5)
45-64	94.9 (93.9-95.8)	83.2 (81.6-84.6)	70.9 (69.3-72.4)
65-74	89.9 (88.4-91.2)	72.9 (71.2-74.6)	55.2 (53.7-56.7)
75+	85.5 (83.4-87.4)	63.6 (61.4-65.6)	44.4 (42.9-46.0)
Male	93.3 (92.4-94.1)	79.7 (78.4-80.9)	66.0 (64.7-67.2)
Female	90.8 (89.6-91.9)	76.5 (75.0-77.9)	61.6 (60.2-63.0)
Diabetes	90.9 (89.4-92.1)	73.3 (71.6-75.0)	53.5 (52.1-54.9)
Hypertension / renal vascular disease	93.3 (91.4-94.8)	79.5 (77.0-81.7)	61.4 (59.1-63.6)
Glomerulonephritis	96.1 (94.4-97.3)	88.7 (86.3-90.7)	82.4 (79.7-84.8)
Other cause	91.6 (90.4-92.6)	78.3 (76.7-79.7)	67.6 (66.1-69.1)
All	92.3 (91.5-92.9)	78.4 (77.4-79.3)	64.2 (63.3-65.1)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival on Dialysis

Table A.5.14

90-day, one- and two-year survival probabilities (cohort 1996-2000): Incident dialysis patients, adjusted from day 1 to 90 days, one and two years, adjusted for age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES			
0-19	99.0 (98.2 - 99.7)	95.9 (94.3 - 97.6)	94.4 (92.3 - 96.5)
20-44	98.5 (98.3 - 98.8)	94.6 (94.0 - 95.1)	89.8 (89.0 - 90.6)
45-64	96.0 (95.8 - 96.3)	87.7 (87.2 - 88.2)	77.7 (77.1 - 78.3)
65-74	91.4 (91.0 - 91.8)	77.0 (76.3 - 77.6)	62.2 (61.4 - 62.9)
75+	87.1 (86.5 - 87.7)	67.4 (66.5 - 68.4)	49.5 (48.5 - 50.5)
Male	94.7 (94.5 - 94.9)	84.9 (84.6 - 85.3)	73.9 (73.4 - 74.4)
Female	94.7 (94.4 - 95.0)	85.1 (84.7 - 85.6)	74.8 (74.2 - 75.4)
Diabetes	94.5 (94.2 - 94.9)	81.9 (81.2 - 82.6)	66.0 (65.2 - 66.9)
Hypertension / renal vascular disease	94.6 (94.2 - 95.0)	84.7 (84.0 - 85.3)	74.0 (73.1 - 74.9)
Glomerulonephritis	96.4 (96.0 - 96.8)	90.3 (89.7 - 91.0)	82.4 (81.5 - 83.3)
Other cause	94.1 (93.8 - 94.3)	84.3 (83.8 - 84.7)	74.4 (73.9 - 75.0)
All	94.7 (94.5 - 94.9)	85.0 (84.7 - 85.3)	74.2 (73.8 - 74.6)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	97.8 (96.7-98.9)	94.5 (92.7-96.3)	91.3 (89.0-93.7)
45-64	95.2 (94.3-96.2)	84.1 (82.4-85.8)	72.3 (70.2-74.5)
65-74	90.8 (89.4-92.2)	75.4 (73.3-77.6)	58.6 (56.1-61.2)
75+	86.2 (84.0-88.5)	65.6 (62.5-68.8)	46.9 (43.7-50.5)
Male	94.5 (93.7-95.2)	83.2 (81.9-84.5)	70.9 (69.3-72.6)
Female	93.9 (93.0-94.8)	83.7 (82.3-85.2)	72.2 (70.4-74.0)
Diabetes	92.9 (91.7-94.0)	78.7 (76.8-80.6)	61.3 (58.9-63.7)
Hypertension / renal vascular disease	95.5 (94.3-96.7)	86.0 (83.9-88.2)	72.1 (69.2-75.0)
Glomerulonephritis	95.6 (93.9-97.2)	87.0 (84.2-89.8)	79.7 (76.3-83.3)
Other cause	92.9 (91.9-94.0)	81.5 (79.8-83.1)	72.0 (70.0-74.0)
All	94.2 (93.6-94.8)	83.4 (82.4-84.3)	71.4 (70.2-72.7)

Table A.5.15

One- and two-year survival probabilities (cohort 1996-2000): Incident dialysis patients, unadjusted
from day 91 to one and two years + 90 days, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	97.0 (95.3 - 98.1)	95.5 (93.3 - 97.0)
20-44	95.0 (94.5 - 95.5)	90.0 (89.2 - 90.7)
45-64	88.1 (87.7 - 88.5)	77.7 (77.2 - 78.3)
65-74	79.2 (78.7 - 79.8)	62.8 (62.3 - 63.3)
75+	71.6 (71.0 - 72.3)	51.9 (51.4 - 52.4)
Male	83.4 (83.0 - 83.7)	69.4 (69.0 - 69.7)
Female	83.1 (82.6 - 83.5)	69.7 (69.3 - 70.2)
Diabetes	79.4 (78.7 - 80.0)	61.3 (60.7 - 61.9)
Hypertension / renal vascular disease	79.9 (79.1 - 80.6)	64.1 (63.3 - 64.8)
Glomerulonephritis	91.5 (90.9 - 92.1)	82.7 (81.9 - 83.5)
Other cause	83.7 (83.3 - 84.1)	71.4 (70.9 - 71.8)
All	83.3 (83.0 - 83.5)	69.5 (69.2 - 69.8)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19	100	100
20-44	96.0 (94.2-97.3)	91.9 (89.5-93.9)
45-64	84.3 (82.7-85.7)	71.7 (70.0-73.3)
65-74	76.2 (74.3-78.0)	57.0 (55.3-58.6)
75+	68.4 (65.9-70.7)	45.6 (43.8-47.3)
Male	81.7 (80.4-82.9)	66.6 (65.2-67.8)
Female	80.1 (78.5-81.6)	64.2 (62.6-65.7)
Diabetes	75.1 (73.3-76.8)	54.4 (52.8-55.9)
Hypertension / renal vascular disease	80.8 (78.2-83.1)	61.4 (59.0-63.7)
Glomerulonephritis	91.0 (88.6-92.9)	83.0 (80.2-85.5)
Other cause	82.2 (80.6-83.7)	70.1 (68.5-71.7)
All	81.1 (80.1-82.0)	65.6 (64.6-66.6)

Table A.5.16

One- and two-year survival probabilities (cohort 1996-2000): Incident dialysis patients, adjusted from day 91 to one and two years + 90 days, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	96.6 (95.1 - 98.2)	94.9 (92.8 - 97.1)
20-44	95.0 (94.4 - 95.5)	89.8 (89.0 - 90.6)
45-64	88.6 (88.1 - 89.1)	78.7 (78.1 - 79.4)
65-74	80.2 (79.6 - 80.8)	64.5 (63.8 - 65.3)
75+	71.9 (70.9 - 72.8)	52.6 (51.6 - 53.7)
Male	86.6 (86.3 - 87.0)	75.1 (74.6 - 75.6)
Female	86.9 (86.5 - 87.4)	76.3 (75.7 - 76.9)
Diabetes	82.1 (81.4 - 82.8)	66.1 (65.2 - 67.0)
Hypertension / renal vascular disease	86.5 (85.8 - 87.1)	75.0 (74.1 - 75.9)
Glomerulonephritis	91.7 (91.0 - 92.3)	83.4 (82.5 - 84.4)
Other cause	86.7 (86.3 - 87.1)	76.4 (75.8 - 77.0)
All	86.8 (86.5 - 87.1)	75.6 (75.1 - 76.0)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	95.9 (94.3-97.5)	91.5 (89.0-94.0)
45-64	85.4 (83.7-87.1)	73.7 (71.5-75.9)
65-74	78.7 (76.5-80.9)	61.1 (58.4-63.8)
75+	70.3 (67.1-73.7)	48.9 (45.3-52.7)
Male	84.7 (83.4-86.1)	71.7 (69.9-73.4)
Female	86.2 (84.9-87.7)	74.1 (72.3-76.1)
Diabetes	79.7 (77.7-81.7)	62.2 (59.7-64.8)
Hypertension / renal vascular disease	86.7 (84.5-88.9)	71.9 (68.9-75.1)
Glomerulonephritis	89.5 (86.9-92.3)	80.6 (77.1-84.3)
Other cause	84.6 (82.9-86.2)	73.7 (71.6-75.8)
All	85.4 (84.4-86.3)	72.7 (71.4-74.0)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival after First Transplant (cadaver donor)

Table A.5.17

One-, two- and five-year survival probabilities (cohort 1993-1997): First transplant patients (cadaver donor), unadjusted from day of transplant to one, two and five years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	96.9 (95.2 - 98.0)	96.1 (94.2 - 97.3)	94.2 (92.1 - 95.7)
20-44	96.9 (96.3 - 97.4)	95.4 (94.7 - 96.0)	90.8 (89.9 - 91.7)
45-64	92.6 (91.8 - 93.3)	89.6 (88.7 - 90.4)	78.6 (77.5 - 79.5)
65+	85.0 (82.0 - 87.6)	80.6 (77.5 - 83.3)	61.6 (58.8 - 64.3)
Male	94.2 (93.6 - 94.7)	91.7 (91.0 - 92.3)	82.8 (81.9 - 83.6)
Female	94.4 (93.7 - 95.1)	92.3 (91.5 - 93.1)	85.4 (84.3 - 86.4)
Diabetes	91.4 (89.6 - 92.8)	87.0 (85.1 - 88.7)	75.0 (73.0 - 77.0)
Hypertension / renal vascular disease	93.2 (91.2 - 94.8)	90.3 (88.1 - 92.1)	78.3 (75.8 - 80.7)
Glomerulonephritis	96.2 (95.4 - 96.8)	94.4 (93.5 - 95.2)	87.4 (86.2 - 88.5)
Other cause	94.1 (93.4 - 94.7)	91.9 (91.2 - 92.6)	84.6 (83.7 - 85.5)
All	94.3 (93.8 - 94.7)	91.9 (91.4 - 92.5)	83.8 (83.1 - 84.4)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19	94.3 (86.2-97.7)	94.3 (86.2-97.7)	92.9 (84.6-96.8)
20-44	98.0 (96.3-98.9)	96.3 (94.3-97.6)	92.4 (89.9-94.3)
45-64	92.5 (90.2-94.3)	88.6 (86.0-90.7)	76.8 (74.0-79.4)
65+	88.7 (79.4-94.0)	83.9 (74.4-90.1)	71.0 (62.1-78.1)
Male	94.6 (92.8-96.0)	91.3 (89.2-93.0)	82.1 (79.7-84.3)
Female	94.8 (92.4-96.4)	92.8 (90.2-94.7)	86.7 (83.7-89.1)
Diabetes	94.4 (89.9-96.9)	89.4 (84.3-93.0)	75.2 (69.7-79.8)
Hypertension / renal vascular disease	89.9 (81.2-94.7)	84.1 (75.2-90.0)	73.8 (65.2-80.6)
Glomerulonephritis	97.5 (95.3-98.7)	95.5 (92.9-97.2)	89.6 (86.3-92.1)
Other cause	93.6 (91.5-95.2)	91.3 (89.0-93.1)	84.0 (81.4-86.3)
All	94.7 (93.3-95.8)	91.9 (90.3-93.2)	83.9 (82.0-85.6)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival after First Transplant (cadaver donor)

Table A.5.18

One-, two- and five-year survival probabilities (cohort 1993-1997): First transplant patients (cadaver donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	96.7 (95.2 - 98.2)	95.8 (94.1 - 97.5)	93.6 (91.5 - 95.7)
20-44	97.2 (96.6 - 97.7)	95.8 (95.1 - 96.4)	91.4 (90.5 - 92.4)
45-64	92.8 (92.1 - 93.6)	89.9 (89.0 - 90.8)	79.1 (77.8 - 80.3)
65+	85.5 (82.3 - 88.7)	81.2 (77.7 - 84.8)	62.4 (58.1 - 67.0)
Male	95.3 (94.7 - 95.8)	93.2 (92.6 - 93.8)	85.4 (84.4 - 86.4)
Female	95.3 (94.6 - 96.0)	93.5 (92.7 - 94.3)	87.3 (86.2 - 88.4)
Diabetes	91.8 (90.1 - 93.4)	87.6 (85.6 - 89.6)	75.6 (73.0 - 78.3)
Hypertension / renal vascular disease	95.7 (94.5 - 96.9)	93.8 (92.3 - 95.3)	85.4 (83.2 - 87.7)
Glomerulonephritis	96.6 (95.9 - 97.3)	95.0 (94.2 - 95.8)	88.5 (87.3 - 89.7)
Other cause	95.0 (94.3 - 95.6)	93.1 (92.4 - 93.8)	86.5 (85.5 - 87.5)
All	95.3 (94.9 - 95.7)	93.3 (92.8 - 93.9)	86.2 (85.4 - 86.9)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	98.1 (96.9-99.3)	96.5 (95.0-98.1)	92.8 (90.6-95.1)
45-64	92.9 (90.9-95.0)	89.2 (86.7-91.7)	77.8 (74.5-81.3)
65+			
Male	95.9 (94.7-97.2)	93.3 (91.7-94.9)	85.7 (83.4-88.1)
Female	95.6 (93.9-97.3)	93.9 (91.9-96.0)	88.6 (85.9-91.3)
Diabetes	95.5 (92.7-98.4)	91.3 (87.5-95.4)	79.0 (73.4-85.0)
Hypertension / renal vascular disease			
Glomerulonephritis	97.5 (95.9-99.1)	95.5 (93.4-97.7)	89.4 (86.2-92.7)
Other cause	94.9 (93.4-96.5)	93.0 (91.2-94.9)	86.9 (84.4-89.4)
All	95.8 (94.8-96.9)	93.6 (92.3-94.9)	86.9 (85.1-88.7)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival after First Transplant (cadaver donor)

Table A.5.19

One- and two-year survival probabilities (cohort 1996-2000): First transplant patients (cadaver donor), unadjusted from day of transplant to one and two years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	98.0 (96.5 - 98.9)	96.7 (94.9 - 97.9)
20-44	97.4 (96.9 - 97.8)	96.2 (95.6 - 96.7)
45-64	94.9 (94.3 - 95.5)	92.4 (91.7 - 93.1)
65+	87.3 (84.9 - 89.4)	82.3 (79.7 - 84.6)
Male	95.3 (94.8 - 95.8)	93.1 (92.5 - 93.7)
Female	96.1 (95.4 - 96.6)	94.1 (93.4 - 94.8)
Diabetes	93.3 (91.8 - 94.4)	90.5 (88.9 - 91.9)
Hypertension / renal vascular disease	93.4 (91.6 - 94.8)	89.7 (87.6 - 91.4)
Glomerulonephritis	96.8 (96.1 - 97.4)	94.8 (94.0 - 95.6)
Other cause	95.9 (95.3 - 96.4)	94.1 (93.5 - 94.7)
All	95.6 (95.2 - 96.0)	93.5 (93.0 - 93.9)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19	98.3 (89.1-99.8)	93.3 (84.2-97.3)
20-44	97.0 (95.2-98.1)	94.9 (92.8-96.4)
45-64	92.5 (90.2-94.3)	88.9 (86.4-91.0)
65+	81.2 (73.3-86.9)	76.5 (68.7-82.6)
Male	94.0 (92.2-95.4)	91.1 (89.0-92.8)
Female	93.8 (91.4-95.5)	90.4 (87.7-92.5)
Diabetes	93.9 (89.9-96.4)	89.8 (85.3-93.0)
Hypertension / renal vascular disease	88.0 (80.7-92.7)	83.7 (76.2-89.0)
Glomerulonephritis	96.3 (93.8-97.8)	93.4 (90.5-95.5)
Other cause	93.4 (91.3-95.1)	90.7 (88.4-92.6)
All	93.9 (92.5-95.0)	90.8 (89.2-92.2)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival after First Transplant (cadaver donor)

Table A.5.20

One- and two-year survival probabilities (cohort 1996-2000): First transplant patients (cadaver donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19		
20-44	97.6 (97.1 - 98.0)	96.4 (95.9 - 97.0)
45-64	95.1 (94.5 - 95.7)	92.7 (92.0 - 93.5)
65+	87.8 (85.4 - 90.3)	83.2 (80.4 - 86.1)
Male	96.3 (95.9 - 96.8)	94.5 (94.0 - 95.1)
Female	96.7 (96.1 - 97.2)	95.1 (94.4 - 95.7)
Diabetes	93.7 (92.4 - 95.0)	91.2 (89.7 - 92.8)
Hypertension / renal vascular disease	95.7 (94.6 - 96.9)	93.4 (91.9 - 94.8)
Glomerulonephritis	97.2 (96.6 - 97.8)	95.5 (94.7 - 96.2)
Other cause	96.5 (96.0 - 97.0)	95.0 (94.4 - 95.6)
All	96.5 (96.1 - 96.8)	94.8 (94.3 - 95.2)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	97.2 (95.9-98.6)	95.3 (93.5-97.0)
45-64	92.8 (90.8-94.9)	89.7 (87.2-92.1)
65+		
Male	95.4 (94.1-96.7)	93.1 (91.5-94.7)
Female	94.6 (92.8-96.5)	91.9 (89.7-94.2)
Diabetes	94.2 (91.2-97.3)	91.0 (87.3-94.8)
Hypertension / renal vascular disease		
Glomerulonephritis	96.4 (94.6-98.4)	93.6 (91.2-96.2)
Other cause	94.6 (93.0-96.3)	92.4 (90.4-94.3)
All	95.1 (94.0-96.2)	92.7 (91.4-94.0)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival after First Transplant (living donor)

Table A.5.21

One-, two- and five-year survival probabilities (cohort 1993-1997): First transplant patients (living donor), unadjusted from day of transplant to one, two and five years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	98.7 (96.6 - 99.5)	98.0 (95.7 - 99.1)	97.4 (94.9 - 98.7)
20-44	98.8 (97.9 - 99.3)	97.7 (96.6 - 98.4)	94.0 (92.5 - 95.2)
45-64	95.1 (92.9 - 96.6)	94.7 (92.5 - 96.2)	88.5 (85.8 - 90.7)
65+	88.5 (78.0 - 94.1)	82.7 (72.1 - 89.5)	69.2 (59.6 - 77.0)
Male	97.6 (96.6 - 98.3)	97.0 (95.9 - 97.8)	93.0 (91.5 - 94.2)
Female	97.4 (96.1 - 98.3)	95.8 (94.2 - 97.0)	91.5 (89.4 - 93.2)
Diabetes	98.2 (94.5 - 99.4)	95.7 (91.5 - 97.9)	83.4 (78.0 - 87.6)
Hypertension / renal vascular disease	89.7 (83.2 - 93.8)	86.9 (80.2 - 91.5)	81.3 (74.4 - 86.5)
Glomerulonephritis	98.7 (97.5 - 99.3)	97.9 (96.5 - 98.7)	95.2 (93.4 - 96.5)
Other cause	97.5 (96.4 - 98.3)	96.8 (95.6 - 97.7)	93.2 (91.6 - 94.5)
All	97.5 (96.8 - 98.1)	96.5 (95.7 - 97.2)	92.4 (91.2 - 93.4)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	100	95.0 (82.6-98.6)	95.0 (82.6-98.6)
45-64			
65+			
Male	97.4 (83.9-99.6)	94.9 (82.2-98.6)	94.9 (82.2-98.6)
Female			
Diabetes			
Hypertension / renal vascular disease			
Glomerulonephritis			
Other cause	96.9 (80.9-99.5)	96.9 (80.9-99.5)	93.8 (78.9-98.3)
All	98.5 (90.2-99.8)	95.5 (87.4-98.5)	94.0 (85.7-97.6)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival after First Transplant (living donor)

Table A.5.22

One-, two- and five-year survival probabilities (cohort 1993-1997): First transplant patients (living donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19			
20-44	98.8 (98.2 - 99.5)	97.7 (96.8 - 98.6)	94.1 (92.6 - 95.5)
45-64	95.2 (93.4 - 97.1)	94.8 (92.9 - 96.7)	88.8 (86.1 - 91.6)
65+			
Male	97.2 (96.2 - 98.2)	96.5 (95.4 - 97.7)	91.7 (90.0 - 93.5)
Female	96.9 (95.5 - 98.3)	94.8 (93.1 - 96.6)	89.3 (86.8 - 91.9)
Diabetes	97.9 (95.7 - 100.0)	95.2 (91.8 - 98.7)	81.2 (75.1 - 87.9)
Hypertension / renal vascular disease			
Glomerulonephritis	98.3 (97.3 - 99.4)	97.4 (96.1 - 98.8)	94.1 (92.1 - 96.1)
Other cause	96.7 (95.5 - 98.0)	95.8 (94.4 - 97.2)	91.0 (89.0 - 93.1)
All	97.1 (96.3 - 97.9)	95.9 (94.9 - 96.9)	90.8 (89.4 - 92.3)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44			
45-64			
65+			
Male			
Female			
Diabetes			
Hypertension / renal vascular disease			
Glomerulonephritis			
Other cause			
All			

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival after First Transplant (living donor)

Table A.5.23

One- and two-year survival probabilities (cohort 1996-2000): First transplant patients (living donor), unadjusted from day of transplant to one and two years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	98.3 (96.4 - 99.2)	98.1 (96.1 - 99.1)
20-44	98.7 (97.9 - 99.2)	98.3 (97.5 - 98.9)
45-64	95.9 (94.2 - 97.1)	95.2 (93.4 - 96.5)
65+	93.9 (85.5 - 97.5)	92.4 (83.8 - 96.6)
Male	98.0 (97.3 - 98.6)	97.6 (96.7 - 98.2)
Female	97.1 (95.8 - 98.0)	96.5 (95.1 - 97.5)
Diabetes	97.4 (94.1 - 98.9)	96.9 (93.5 - 98.6)
Hypertension / renal vascular disease	90.4 (84.7 - 94.0)	89.6 (83.9 - 93.4)
Glomerulonephritis	99.0 (98.0 - 99.5)	98.4 (97.2 - 99.0)
Other cause	97.7 (96.8 - 98.4)	97.4 (96.4 - 98.1)
All	97.7 (97.1 - 98.2)	97.2 (96.5 - 97.8)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	98.8 (91.8-99.8)	97.5 (90.8-99.4)
45-64	100	100
65+		
Male	99.0 (93.0-99.8)	97.9 (92.1-99.5)
Female	98.0 (87.1-99.7)	98.0 (87.1-99.7)
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis	100	98.1 (87.6-99.7)
Other cause	98.8 (91.8-99.8)	98.8 (91.8-99.8)
All	98.6 (94.7-99.6)	97.9 (93.9-99.3)

A5 SURVIVAL PROBABILITY (residents only)
Patient Survival after First Transplant (living donor)

Table A.5.24

One- and two-year survival probabilities (cohort 1996-2000): First transplant patients (living donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19		
20-44	98.7 (98.1 - 99.3)	98.3 (97.7 - 99.0)
45-64	96.1 (94.7 - 97.5)	95.4 (93.9 - 97.0)
65+		
Male	97.8 (97.1 - 98.6)	97.3 (96.5 - 98.2)
Female	96.6 (95.3 - 97.9)	95.9 (94.5 - 97.4)
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis		
Other cause	97.2 (96.3 - 98.2)	96.8 (95.7 - 97.8)
All	97.4 (96.7 - 98.1)	96.8 (96.1 - 97.6)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44		
45-64		
65+		
Male		
Female		
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis		
Other cause		
All		

A5 SURVIVAL PROBABILITY (residents only)
Graft Survival after First Transplant (cadaver donor)

Table A.5.25

One-, two- and five-year survival probabilities (cohort 1993-1997): First graft (cadaver donor), unadjusted
from day of transplant to one, two and five years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	83.4 (80.5-85.8)	79.4 (76.5-82.0)	67.8 (65.0-70.4)
20-44	88.0 (87.0-89.0)	84.9 (83.8-85.9)	74.8 (73.7-75.9)
45-64	84.4 (83.4-85.3)	81.0 (80.0-81.9)	69.0 (68.0-70.0)
65+	80.8 (77.7-83.5)	76.6 (73.5-79.4)	56.7 (54.1-59.2)
Male	85.5 (84.7-86.3)	82.1 (81.2-82.9)	69.9 (69.0-70.7)
Female	85.7 (84.7-86.7)	82.5 (81.4-83.6)	71.8 (70.6-72.9)
Diabetes	83.5 (81.5-85.3)	79.0 (77.0-80.9)	65.8 (63.8-67.7)
Hypertension / renal vascular disease	84.1 (81.6-86.3)	79.3 (76.8-81.6)	66.0 (63.6-68.3)
Glomerulonephritis	87.9 (86.7-89.0)	84.7 (83.5-85.9)	72.9 (71.6-74.1)
Other cause	85.0 (84.1-85.9)	82.0 (81.1-82.9)	71.2 (70.2-72.1)
All	85.6 (85.0-86.2)	82.2 (81.5-82.9)	70.6 (69.9-71.3)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19	80.6 (71.2-87.3)	75.8 (66.6-82.8)	69.4 (60.6-76.5)
20-44	90.4 (87.7-92.5)	87.7 (84.8-90.0)	77.0 (73.9-79.8)
45-64	86.4 (83.7-88.7)	81.8 (79.0-84.2)	68.3 (65.6-70.9)
65+	85.5 (76.0-91.4)	82.3 (72.8-88.7)	69.4 (60.6-76.5)
Male	87.2 (84.9-89.2)	82.7 (80.3-84.9)	69.6 (67.2-71.9)
Female	88.5 (85.6-90.8)	85.8 (82.8-88.3)	75.8 (72.7-78.7)
Diabetes	87.0 (81.6-90.8)	81.4 (75.9-85.7)	66.5 (61.4-71.1)
Hypertension / renal vascular disease	78.3 (69.4-84.8)	73.9 (65.3-80.7)	60.7 (53.3-67.3)
Glomerulonephritis	89.7 (86.4-92.2)	86.2 (82.8-89.1)	75.4 (71.8-78.6)
Other cause	87.8 (85.3-90.0)	84.4 (81.7-86.7)	72.9 (70.1-75.4)
All	87.7 (85.9-89.3)	83.9 (82.0-85.6)	72.0 (70.1-73.8)

A5 SURVIVAL PROBABILITY (residents only)
Graft Survival after First Transplant (cadaver donor)

Table A.5.26

One-, two- and five-year survival probabilities (cohort 1993-1997): First graft (cadaver donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	82.7 (79.5-86.0)	78.6 (75.1-82.2)	66.7 (62.7-70.9)
20-44	88.4 (87.3-89.4)	85.3 (84.1-86.5)	75.4 (74.0-76.9)
45-64	84.5 (83.4-85.6)	81.1 (79.9-82.3)	69.2 (67.8-70.7)
65+	81.1 (77.6-84.8)	77.0 (73.2-80.9)	57.3 (52.9-62.0)
Male	85.8 (84.9-86.8)	82.4 (81.4-83.4)	70.4 (69.1-71.6)
Female	85.8 (84.5-87.0)	82.5 (81.2-83.9)	71.8 (70.3-73.4)
Diabetes	83.4 (81.1-85.8)	78.9 (76.4-81.5)	65.6 (62.7-68.6)
Hypertension / renal vascular disease	85.4 (82.9-88.0)	81.0 (78.1-83.9)	68.4 (65.1-71.9)
Glomerulonephritis	88.0 (86.7-89.3)	84.8 (83.4-86.2)	72.9 (71.1-74.7)
Other cause	85.0 (84.0-86.1)	82.0 (80.9-83.2)	71.2 (69.8-72.5)
All	85.8 (85.1-86.6)	82.5 (81.7-83.3)	71.0 (70.0-72.0)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	90.7 (88.1-93.3)	88.0 (85.2-90.9)	77.5 (73.9-81.3)
45-64	86.6 (83.8-89.4)	82.0 (78.9-85.2)	68.6 (64.9-72.6)
65+			
Male	87.6 (85.3-90.1)	83.3 (80.6-86.0)	70.4 (67.2-73.8)
Female	88.5 (85.6-91.5)	85.8 (82.7-89.1)	75.8 (71.9-79.8)
Diabetes	87.2 (82.3-92.4)	81.7 (76.0-87.8)	66.9 (60.1-74.5)
Hypertension / renal vascular disease	79.9 (71.3-89.4)	75.8 (66.7-86.1)	63.3 (53.2-75.2)
Glomerulonephritis	89.4 (86.2-92.7)	85.9 (82.3-89.7)	74.8 (70.3-79.5)
Other cause	88.1 (85.5-90.7)	84.7 (81.8-87.6)	73.3 (69.8-76.9)
All	88.0 (86.2-89.9)	84.3 (82.2-86.4)	72.5 (70.0-75.1)

A5 SURVIVAL PROBABILITY (residents only)
Graft Survival after First Transplant (cadaver donor)

Table A.5.27

One- and two-year survival probabilities (cohort 1996-2000): First graft (cadaver donor), unadjusted
from day of transplant to one and two years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	88.8 (86.1-91.1)	85.1 (82.2-87.6)
20-44	90.5 (89.6-91.3)	88.2 (87.3-89.1)
45-64	88.9 (88.1-89.7)	85.4 (84.5-86.2)
65+	82.9 (80.3-85.2)	77.0 (74.4-79.4)
Male	89.2 (88.5-89.9)	85.8 (85.1-86.6)
Female	89.0 (88.0-89.8)	86.1 (85.1-87.0)
Diabetes	88.0 (86.3-89.5)	84.5 (82.7-86.1)
Hypertension / renal vascular disease	85.5 (83.2-87.4)	81.1 (78.8-83.2)
Glomerulonephritis	90.1 (89.0-91.1)	87.0 (85.8-88.1)
Other cause	89.5 (88.7-90.2)	86.5 (85.6-87.3)
All	89.1 (88.6-89.7)	85.9 (85.3-86.5)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19	84.9 (74.5-91.3)	79.2 (68.9-86.5)
20-44	91.4 (88.9-93.4)	88.5 (85.8-90.7)
45-64	88.6 (86.0-90.7)	84.1 (81.4-86.5)
65+	78.8 (71.0-84.8)	74.1 (66.4-80.3)
Male	89.1 (86.9-90.9)	84.9 (82.5-86.9)
Female	88.7 (85.9-91.0)	85.4 (82.4-87.9)
Diabetes	89.3 (84.8-92.6)	84.3 (79.4-88.1)
Hypertension / renal vascular disease	78.3 (70.8-84.0)	73.9 (66.6-79.9)
Glomerulonephritis	91.0 (87.8-93.4)	86.9 (83.4-89.7)
Other cause	89.3 (86.8-91.3)	86.0 (83.4-88.3)
All	88.9 (87.3-90.4)	85.1 (83.3-86.7)

A5 SURVIVAL PROBABILITY (residents only)
Graft Survival after First Transplant (cadaver donor)

Table A.5.28

One- and two-year survival probabilities (cohort 1996-2000): First graft (cadaver donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	88.3 (85.4-91.3)	84.4 (81.1-87.8)
20-44	90.7 (89.7-91.6)	88.4 (87.4-89.4)
45-64	89.0 (88.1-89.9)	85.6 (84.6-86.6)
65+	83.3 (80.4-86.2)	77.5 (74.4-80.8)
Male	89.6 (88.9-90.4)	86.4 (85.5-87.2)
Female	89.1 (88.0-90.1)	86.2 (85.1-87.3)
Diabetes	88.0 (86.2-89.8)	84.5 (82.5-86.5)
Hypertension / renal vascular disease	86.7 (84.5-89.1)	82.7 (80.2-85.3)
Glomerulonephritis	90.2 (89.0-91.3)	87.1 (85.8-88.4)
Other cause	89.6 (88.8-90.4)	86.6 (85.7-87.6)
All	89.4 (88.8-90.0)	86.3 (85.6-87.0)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	91.7 (89.4-94.1)	88.9 (86.2-91.6)
45-64	88.7 (86.1-91.3)	84.5 (81.5-87.5)
65+		
Male	89.6 (87.5-91.8)	85.6 (83.1-88.1)
Female	88.7 (85.9-91.6)	85.5 (82.4-88.7)
Diabetes	88.9 (84.7-93.4)	84.4 (79.5-89.6)
Hypertension / renal vascular disease	80.0 (72.6-88.2)	75.9 (68.0-84.8)
Glomerulonephritis	90.8 (87.8-93.9)	86.6 (83.1-90.3)
Other cause	89.5 (87.1-92.0)	86.3 (83.6-89.1)
All	89.2 (87.5-91.0)	85.5 (83.6-87.5)

A5 SURVIVAL PROBABILITY (residents only)
Graft Survival after First Transplant (living donor)

Table A.5.29

One-, two- and five-year survival probabilities (cohort 1993-1997): First graft (living donor), unadjusted
from day of transplant to one, two and five years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	93.0 (89.6-95.4)	91.6 (88.0-94.1)	82.3 (78.2-85.7)
20-44	93.3 (91.7-94.6)	90.4 (88.6-91.9)	82.6 (80.6-84.4)
45-64	90.3 (87.7-92.4)	89.3 (86.6-91.5)	82.1 (79.1-84.7)
65+	88.2 (77.6-94.0)	80.4 (69.8-87.6)	66.7 (57.2-74.5)
Male	93.0 (91.5-94.2)	90.7 (89.1-92.1)	82.4 (80.5-84.1)
Female	91.2 (89.1-93.0)	88.8 (86.5-90.7)	81.3 (78.8-83.5)
Diabetes	93.7 (89.0-96.4)	89.3 (84.1-92.9)	76.7 (71.2-81.3)
Hypertension / renal vascular disease	85.6 (78.6-90.4)	79.8 (72.8-85.2)	70.2 (63.5-75.9)
Glomerulonephritis	94.5 (92.6-95.9)	92.8 (90.6-94.4)	85.2 (82.6-87.4)
Other cause	91.4 (89.6-92.9)	89.4 (87.5-91.0)	82.0 (79.9-83.9)
All	92.3 (91.1-93.4)	90.0 (88.7-91.2)	82.0 (80.5-83.4)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	91.4 (77.7-96.9)	85.7 (72.1-93.0)	77.1 (64.2-85.9)
45-64			
65+			
Male	88.2 (74.3-94.9)	85.3 (71.4-92.8)	79.4 (66.0-88.0)
Female			
Diabetes			
Hypertension / renal vascular disease			
Glomerulonephritis			
Other cause			
All	93.1 (83.7-97.2)	89.7 (79.9-94.8)	81.0 (71.2-87.8)

A5 SURVIVAL PROBABILITY (residents only)
Graft Survival after First Transplant (living donor)

Table A.5.30

One-, two- and five-year survival probabilities (cohort 1993-1997): First graft (living donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	92.7 (89.6-95.9)	91.2 (87.8-94.7)	81.7 (77.0-86.5)
20-44	93.2 (91.7-94.8)	90.3 (88.5-92.1)	82.3 (80.0-84.7)
45-64	90.4 (87.9-93.0)	89.4 (86.8-92.2)	82.2 (78.9-85.6)
65+			
Male	92.4 (90.8-94.1)	90.0 (88.2-91.9)	81.0 (78.6-83.6)
Female	90.6 (88.3-92.9)	87.9 (85.4-90.5)	79.7 (76.6-83.0)
Diabetes	93.5 (89.7-97.5)	89.0 (84.2-94.0)	76.0 (69.6-83.1)
Hypertension / renal vascular disease	85.8 (79.5-92.7)	80.1 (72.9-88.1)	70.5 (62.3-79.8)
Glomerulonephritis	94.2 (92.4-96.1)	92.4 (90.3-94.6)	84.5 (81.6-87.5)
Other cause	90.6 (88.6-92.6)	88.4 (86.2-90.7)	80.3 (77.6-83.2)
All	91.7 (90.4-93.1)	89.2 (87.7-90.8)	80.6 (78.6-82.7)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44			
45-64			
65+			
Male			
Female			
Diabetes			
Hypertension / renal vascular disease			
Glomerulonephritis			
Other cause			
All			

A5 SURVIVAL PROBABILITY (residents only)
Graft Survival after First Transplant (living donor)

Table A.5.31

One- and two-year survival probabilities (cohort 1996-2000): First graft (living donor), unadjusted
from day of transplant to one and two years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	95.2 (92.4-97.0)	93.6 (90.6-95.7)
20-44	94.5 (93.2-95.6)	92.9 (91.4-94.1)
45-64	92.4 (90.3-94.0)	91.1 (89.0-92.8)
65+	93.9 (85.5-97.5)	92.4 (83.8-96.6)
Male	95.0 (93.9-96.0)	93.7 (92.4-94.7)
Female	92.0 (90.2-93.6)	90.2 (88.2-91.9)
Diabetes	93.1 (88.9-95.8)	91.6 (87.1-94.5)
Hypertension / renal vascular disease	87.3 (81.4-91.4)	84.3 (78.3-88.8)
Glomerulonephritis	95.3 (93.6-96.5)	94.0 (92.2-95.5)
Other cause	94.0 (92.7-95.2)	92.5 (91.1-93.8)
All	94.0 (93.0-94.8)	92.5 (91.4-93.4)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	94.4 (86.6-97.8)	93.1 (85.0-96.9)
45-64	97.0 (81.4-99.5)	93.9 (79.5-98.3)
65+		
Male	95.2 (88.2-98.1)	94.0 (86.8-97.3)
Female	95.6 (84.3-98.8)	93.3 (82.0-97.6)
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis	92.7 (80.5-97.4)	92.7 (80.5-97.4)
Other cause	97.3 (90.0-99.3)	94.6 (86.9-97.8)
All	95.3 (90.2-97.8)	93.8 (88.4-96.7)

A5 SURVIVAL PROBABILITY (residents only)
Graft Survival after First Transplant (living donor)

Table A.5.32

One- and two-year survival probabilities (cohort 1996-2000): First graft (living donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	95.0 (92.6-97.5)	93.4 (90.6-96.2)
20-44	94.4 (93.2-95.7)	92.8 (91.3-94.3)
45-64	92.4 (90.5-94.4)	91.1 (89.0-93.3)
65+		
Male	94.8 (93.6-96.0)	93.4 (92.1-94.7)
Female	91.8 (89.9-93.8)	89.9 (87.8-92.1)
Diabetes	93.0 (89.4-96.8)	91.4 (87.5-95.5)
Hypertension / renal vascular disease		
Glomerulonephritis	95.0 (93.4-96.7)	93.7 (91.9-95.6)
Other cause	93.8 (92.4-95.3)	92.3 (90.7-93.9)
All	93.7 (92.7-94.8)	92.1 (90.9-93.3)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44		
45-64		
65+		
Male		
Female		
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis		
Other cause		
All		