



EUROPEAN RENAL ASSOCIATION

EUROPEAN DIALYSIS AND TRANSPLANT ASSOCIATION

ERA - EDTA REGISTRY

Survival data

Austria

2004

ERA-EDTA Registry

Academic Medical Center
Department of Medical Informatics
Room J1B-125
PO Box 22700
1100 DE Amsterdam
The Netherlands

Telephone +31 20 566 7637
Fax +31 20 691 9840
E-mail erareg@amc.uva.nl
Website www.era-edta-reg.org

Further information (e.g. collaborating registries, methods) can be found in the ERA-EDTA Registry annual report 2004, which is published on the ERA-EDTA Registry website www.era-edta-reg.org.

Table of Contents

A5 SURVIVAL PROBABILITY

Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.1	5
90-day, one-, two- and five-year survival probabilities (cohort 1995-1999): Incident RRT patients, unadjusted <i>from day 1 to 90 days, one, two and five years, by age, gender and primary diagnosis</i>	
Table A.5.2	6
90-day, one-, two- and five-year survival probabilities (cohort 1995-1999): Incident RRT patients, adjusted <i>from day 1 to 90 days, one, two and five years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.3	7
One-, two- and five-year survival probabilities (cohort 1995-1999): Incident RRT patients, unadjusted <i>from day 91 to one, two and five years + 90 days, by age, gender and primary diagnosis</i>	
Table A.5.4	8
One-, two- and five-year survival probabilities (cohort 1995-1999): Incident RRT patients, adjusted <i>from day 91 to one, two and five years + 90 days, adjusted for age, gender and primary diagnosis</i>	
Table A.5.5	9
90-day, one- and two-year survival probabilities (cohort 1998-2002): Incident RRT patients, unadjusted <i>from day 1 to 90 days, one and two years, by age, gender and primary diagnosis</i>	
Table A.5.6	10
90-day, one- and two-year survival probabilities (cohort 1998-2002): Incident RRT patients, adjusted <i>from day 1 to 90 days, one and two years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.7	11
One- and two-year survival probabilities (cohort 1998-2002): Incident RRT patients, unadjusted <i>from day 91 to one and two years + 90 days, by age, gender and primary diagnosis</i>	
Table A.5.8	12
One- and two-year survival probabilities (cohort 1998-2002): Incident RRT patients, adjusted <i>from day 91 to one and two years + 90 days, adjusted for age, gender and primary diagnosis</i>	

Patient Survival on Dialysis

Table A.5.9	13
90-day, one-, two- and five-year survival probabilities (cohort 1995-1999): Incident dialysis patients, unadjusted <i>from day 1 to 90 days, one, two and five years, by age, gender and primary diagnosis</i>	
Table A.5.10	14
90-day, one-, two- and five-year survival probabilities (cohort 1995-1999): Incident dialysis patients, adjusted <i>from day 1 to 90 days, one, two and five years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.11	15
One-, two- and five-year survival probabilities (cohort 1995-1999): Incident dialysis patients, unadjusted <i>from day 91 to one, two and five years + 90 days, by age, gender and primary diagnosis</i>	
Table A.5.12	16
One-, two- and five-year survival probabilities (cohort 1995-1999): Incident dialysis patients, adjusted <i>from day 91 to one, two and five years + 90 days, adjusted for age, gender and primary diagnosis</i>	
Table A.5.13	17
90-day, one- and two-year survival probabilities (cohort 1998-2002): Incident dialysis patients, unadjusted <i>from day 1 to 90 days, one and two years, by age, gender and primary diagnosis</i>	
Table A.5.14	18
90-day, one- and two-year survival probabilities (cohort 1998-2002): Incident dialysis patients, adjusted <i>from day 1 to 90 days, one and two years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.15	19
One- and two-year survival probabilities (cohort 1998-2002): Incident dialysis patients, unadjusted <i>from day 91 to one and two years + 90 days, by age, gender and primary diagnosis</i>	
Table A.5.16	20
One- and two-year survival probabilities (cohort 1998-2002): Incident dialysis patients, adjusted <i>from day 91 to one and two years + 90 days, adjusted for age, gender and primary diagnosis</i>	

Patient Survival after First Transplant (cadaver donor)

Table A.5.17	21
One-, two- and five-year survival probabilities (cohort 1995-1999): First transplant patients (cadaver donor), unadjusted <i>from day of transplant to one, two and five years, by age, gender and primary diagnosis</i>	
Table A.5.18	22
One-, two- and five-year survival probabilities (cohort 1995-1999): First transplant patients (cadaver donor), adjusted <i>from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis</i>	

Table A.5.19	23
One- and two-year survival probabilities (cohort 1998-2002): First transplant patients (cadaver donor), unadjusted <i>from day of transplant to one and two years, by age, gender and primary diagnosis</i>	
Table A.5.20	24
One- and two-year survival probabilities (cohort 1998-2002): First transplant patients (cadaver donor), adjusted <i>from day of transplant to one and two years, adjusted for age, gender and primary diagnosis</i>	
Patient Survival after First Transplant (living donor)	
Table A.5.21	25
One-, two- and five-year survival probabilities (cohort 1995-1999): First transplant patients (living donor), unadjusted <i>from day of transplant to one, two and five years, by age, gender and primary diagnosis</i>	
Table A.5.22	26
One-, two- and five-year survival probabilities (cohort 1995-1999): First transplant patients (living donor), adjusted <i>from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.23	27
One- and two-year survival probabilities (cohort 1998-2002): First transplant patients (living donor), unadjusted <i>from day of transplant to one and two years, by age, gender and primary diagnosis</i>	
Table A.5.24	28
One- and two-year survival probabilities (cohort 1998-2002): First transplant patients (living donor), adjusted <i>from day of transplant to one and two years, adjusted for age, gender and primary diagnosis</i>	
Graft Survival after First Transplant (cadaver donor)	
Table A.5.25	29
One-, two- and five-year survival probabilities (cohort 1995-1999): First graft (cadaver donor), unadjusted <i>from day of transplant to one, two and five years, by age, gender and primary diagnosis</i>	
Table A.5.26	30
One-, two- and five-year survival probabilities (cohort 1995-1999): First graft (cadaver donor), adjusted <i>from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.27	31
One- and two-year survival probabilities (cohort 1998-2002): First graft (cadaver donor), unadjusted <i>from day of transplant to one and two years, by age, gender and primary diagnosis</i>	
Table A.5.28	32
One- and two-year survival probabilities (cohort 1998-2002): First graft (cadaver donor), adjusted <i>from day of transplant to one and two years, adjusted for age, gender and primary diagnosis</i>	
Graft Survival after First Transplant (living donor)	
Table A.5.29	33
One-, two- and five-year survival probabilities (cohort 1995-1999): First graft (living donor), unadjusted <i>from day of transplant to one, two and five years, by age, gender and primary diagnosis</i>	
Table A.5.30	34
One-, two- and five-year survival probabilities (cohort 1995-1999): First graft (living donor), adjusted <i>from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis</i>	
Table A.5.31	35
One- and two-year survival probabilities (cohort 1998-2002): First graft (living donor), unadjusted <i>from day of transplant to one and two years, by age, gender and primary diagnosis</i>	
Table A.5.32	36
One- and two-year survival probabilities (cohort 1998-2002): First graft (living donor), adjusted <i>from day of transplant to one and two years, adjusted for age, gender and primary diagnosis</i>	

A5 SURVIVAL PROBABILITY
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.1

90-day, one-, two- and five-year survival probabilities (cohort 1995-1999): Incident RRT patients, unadjusted
from day 1 to 90 days, one, two and five years, by age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES				
0-19	99.2 (98.5-99.6)	97.1 (96.0-97.9)	95.6 (94.3-96.6)	93.0 (91.4-94.2)
20-44	98.7 (98.4-98.9)	95.7 (95.3-96.0)	92.4 (91.9-92.8)	85.0 (84.4-85.6)
45-64	96.4 (96.2-96.7)	88.7 (88.3-89.0)	80.1 (79.7-80.5)	60.1 (59.7-60.5)
65-74	92.3 (92.0-92.6)	78.4 (78.0-78.9)	64.1 (63.7-64.5)	33.3 (33.1-33.6)
75+	88.5 (88.0-89.0)	69.6 (69.0-70.1)	51.9 (51.5-52.4)	18.8 (18.7-19.0)
Men	94.2 (93.9-94.4)	83.3 (83.0-83.6)	72.2 (71.9-72.5)	48.2 (48.0-48.4)
Women	93.9 (93.6-94.2)	82.8 (82.5-83.2)	71.6 (71.3-72.0)	48.7 (48.4-49.0)
Diabetes	94.2 (93.9-94.6)	80.2 (79.7-80.8)	63.4 (62.9-63.9)	32.5 (32.2-32.7)
Hypertension / renal vascular disease	92.6 (92.2-93.1)	79.8 (79.3-80.4)	66.3 (65.7-66.9)	37.3 (37.0-37.6)
Glomerulonephritis	96.8 (96.4-97.1)	91.2 (90.7-91.7)	84.8 (84.2-85.4)	68.0 (67.3-68.6)
Other cause	93.7 (93.4-93.9)	82.9 (82.5-83.2)	73.2 (72.9-73.6)	52.2 (51.9-52.5)
All	94.1 (93.9-94.2)	83.1 (82.9-83.4)	72.0 (71.7-72.2)	48.4 (48.2-48.6)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA				
0-19	100	97.2 (89.6-99.3)	95.8 (88.0-98.5)	91.4 (83.0-95.8)
20-44	97.9 (96.6-98.7)	95.2 (93.4-96.4)	92.5 (90.5-94.1)	83.3 (80.9-85.4)
45-64	94.6 (93.5-95.5)	83.3 (81.8-84.7)	72.5 (71.0-74.0)	48.7 (47.5-49.8)
65-74	90.0 (88.5-91.3)	71.5 (69.7-73.2)	55.3 (53.8-56.7)	22.1 (21.6-22.5)
75+	86.5 (84.3-88.5)	63.7 (61.4-65.9)	42.4 (40.9-44.0)	11.2 (11.0-11.5)
Men	93.4 (92.5-94.2)	80.0 (78.8-81.2)	67.6 (66.4-68.7)	42.1 (41.3-42.9)
Women	91.3 (90.1-92.4)	77.0 (75.5-78.4)	64.3 (62.9-65.6)	39.6 (38.7-40.5)
Diabetes	91.3 (89.9-92.6)	73.9 (72.2-75.6)	55.5 (54.1-57.0)	25.3 (24.7-25.9)
Hypertension / renal vascular disease	93.3 (91.2-94.9)	78.5 (75.9-80.9)	61.8 (59.4-64.1)	32.1 (31.0-33.3)
Glomerulonephritis	96.4 (94.7-97.5)	89.6 (87.3-91.4)	83.8 (81.3-86.0)	67.4 (65.0-69.7)
Other cause	91.9 (90.7-92.9)	78.6 (77.1-80.0)	69.2 (67.8-70.6)	46.2 (45.1-47.2)
All	92.5 (91.8-93.2)	78.8 (77.8-79.7)	66.2 (65.3-67.1)	41.1 (40.5-41.7)

A5 SURVIVAL PROBABILITY
Patient Survival on Renal Replacement Therapy (RRT)

Table A.5.2

90-day, one-, two- and five-year survival probabilities (cohort 1995-1999): Incident RRT patients, adjusted from day 1 to 90 days, one, two and five years, adjusted for age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES				
0-19	99.1 (98.5-99.7)	96.7 (95.5-97.8)	95.0 (93.6-96.4)	91.9 (90.2-93.7)
20-44	98.6 (98.4-98.8)	95.5 (95.1-95.9)	92.1 (91.6-92.7)	84.4 (83.7-85.2)
45-64	96.5 (96.3-96.8)	89.0 (88.6-89.4)	80.6 (80.1-81.1)	60.4 (59.8-61.0)
65-74	92.6 (92.3-93.0)	79.4 (78.8-79.9)	65.4 (64.7-66.0)	34.3 (33.6-34.9)
75+	88.5 (87.9-89.0)	69.7 (69.0-70.5)	52.1 (51.2-53.0)	18.6 (17.9-19.3)
Men	95.8 (95.6-95.9)	87.5 (87.2-87.8)	78.1 (77.8-78.5)	53.6 (53.1-54.1)
Women	95.9 (95.7-96.1)	88.0 (87.6-88.3)	79.1 (78.6-79.5)	56.5 (55.9-57.1)
Diabetes	95.5 (95.2-95.8)	84.1 (83.5-84.7)	69.2 (68.5-70.0)	36.7 (35.9-37.6)
Hypertension / renal vascular disease	95.9 (95.6-96.2)	88.3 (87.8-88.7)	79.2 (78.6-79.9)	54.4 (53.6-55.3)
Glomerulonephritis	97.0 (96.7-97.3)	91.5 (91.0-92.0)	85.0 (84.4-85.7)	65.5 (64.5-66.5)
Other cause	95.4 (95.2-95.6)	87.0 (86.7-87.4)	79.0 (78.5-79.4)	58.0 (57.5-58.6)
All	95.8 (95.7-95.9)	87.7 (87.4-87.9)	78.5 (78.2-78.8)	54.8 (54.4-55.2)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA				
0-19				
20-44	97.9 (96.8-98.9)	95.0 (93.4-96.6)	92.2 (90.3-94.3)	82.7 (79.9-85.6)
45-64	95.0 (94.0-96.0)	84.5 (82.9-86.1)	74.3 (72.3-76.3)	50.6 (48.3-53.0)
65-74	91.0 (89.6-92.4)	74.7 (72.6-76.9)	59.6 (57.2-62.2)	25.8 (23.6-28.3)
75+	87.6 (85.3-89.9)	66.4 (63.1-69.7)	45.6 (42.2-49.3)	13.4 (11.0-16.2)
Men	95.1 (94.5-95.8)	84.7 (83.5-85.9)	73.5 (72.0-75.1)	45.5 (43.6-47.5)
Women	94.8 (94.0-95.5)	85.3 (84.0-86.6)	75.4 (73.8-77.1)	49.5 (47.3-51.8)
Diabetes	93.7 (92.6-94.8)	80.4 (78.6-82.2)	64.3 (62.0-66.6)	31.7 (29.2-34.3)
Hypertension / renal vascular disease	96.1 (94.9-97.2)	86.7 (84.7-88.8)	74.4 (71.6-77.3)	45.4 (41.8-49.3)
Glomerulonephritis	95.8 (94.2-97.4)	87.5 (84.8-90.3)	80.1 (76.8-83.5)	57.5 (53.3-62.0)
Other cause	93.7 (92.8-94.7)	82.8 (81.3-84.4)	74.4 (72.6-76.2)	49.2 (46.9-51.5)
All	94.9 (94.4-95.5)	84.9 (84.0-85.8)	74.3 (73.2-75.4)	47.1 (45.6-48.6)

Table A.5.3

One-, two- and five-year survival probabilities (cohort 1995-1999): Incident RRT patients, unadjusted
from day 91 to one, two and five years + 90 days, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	97.4 (96.3-98.3)	96.1 (94.8-97.1)	93.3 (91.6-94.6)
20-44	96.1 (95.7-96.4)	92.7 (92.2-93.2)	85.1 (84.5-85.7)
45-64	89.4 (89.1-89.8)	81.0 (80.6-81.4)	60.4 (60.0-60.8)
65-74	81.0 (80.5-81.4)	65.9 (65.5-66.4)	33.8 (33.6-34.0)
75+	73.4 (72.8-74.0)	54.4 (53.9-54.9)	19.3 (19.2-19.5)
Men	85.3 (85.0-85.6)	73.8 (73.5-74.1)	49.0 (48.7-49.2)
Women	84.9 (84.5-85.3)	73.6 (73.2-74.0)	49.7 (49.4-50.0)
Diabetes	80.4 (79.8-80.9)	63.6 (63.0-64.1)	31.9 (31.7-32.2)
Hypertension / renal vascular disease	82.3 (81.7-82.9)	68.1 (67.5-68.7)	38.2 (37.8-38.5)
Glomerulonephritis	92.4 (91.9-92.9)	86.0 (85.4-86.6)	68.3 (67.7-68.9)
Other cause	85.7 (85.3-86.0)	75.6 (75.3-76.0)	53.7 (53.4-54.0)
All	85.2 (84.9-85.4)	73.7 (73.5-74.0)	49.3 (49.1-49.5)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19	96.0 (85.7-98.9)	96.0 (85.7-98.9)	91.8 (81.1-96.6)
20-44	96.1 (94.4-97.3)	93.3 (91.3-94.9)	82.8 (80.2-85.1)
45-64	85.3 (83.7-86.7)	73.9 (72.3-75.4)	49.5 (48.3-50.8)
65-74	74.7 (72.8-76.5)	57.1 (55.5-58.7)	22.7 (22.2-23.2)
75+	66.9 (64.4-69.3)	42.3 (40.6-43.9)	11.2 (10.9-11.5)
Men	81.8 (80.5-83.0)	68.5 (67.2-69.7)	42.4 (41.6-43.2)
Women	80.6 (79.0-82.1)	66.5 (65.0-68.0)	40.9 (39.9-41.9)
Diabetes	75.7 (73.8-77.4)	56.5 (54.9-58.0)	25.1 (24.5-25.7)
Hypertension / renal vascular disease	79.7 (76.9-82.1)	62.0 (59.5-64.4)	32.9 (31.6-34.1)
Glomerulonephritis	91.6 (89.4-93.4)	84.9 (82.3-87.1)	67.5 (65.0-70.0)
Other cause	82.2 (80.6-83.6)	71.5 (69.9-73.0)	47.7 (46.6-48.9)
All	81.3 (80.3-82.2)	67.7 (66.7-68.6)	41.8 (41.2-42.4)

Table A.5.4

One-, two- and five-year survival probabilities (cohort 1995-1999): Incident RRT patients, adjusted from day 91 to one, two and five years + 90 days, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	97.0 (95.9-98.2)	95.5 (94.1-96.9)	92.2 (90.4-94.0)
20-44	96.0 (95.6-96.4)	92.5 (92.0-93.1)	84.7 (84.0-85.5)
45-64	89.8 (89.4-90.2)	81.5 (81.0-82.0)	61.1 (60.5-61.8)
65-74	81.9 (81.3-82.4)	67.2 (66.5-67.9)	35.2 (34.5-35.9)
75+	73.6 (72.8-74.4)	54.5 (53.6-55.4)	19.4 (18.6-20.1)
Men	88.9 (88.6-89.2)	79.2 (78.9-79.6)	54.3 (53.8-54.9)
Women	89.3 (89.0-89.7)	80.3 (79.8-80.7)	57.3 (56.6-57.9)
Diabetes	84.1 (83.5-84.7)	69.1 (68.4-69.9)	36.3 (35.4-37.2)
Hypertension / renal vascular disease	89.6 (89.2-90.1)	80.2 (79.6-80.8)	55.2 (54.3-56.1)
Glomerulonephritis	92.6 (92.1-93.1)	86.0 (85.3-86.7)	66.0 (65.0-67.0)
Other cause	89.1 (88.7-89.4)	80.6 (80.2-81.0)	59.3 (58.7-59.9)
All	89.1 (88.9-89.3)	79.7 (79.4-80.0)	55.5 (55.1-56.0)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	96.0 (94.5-97.5)	93.2 (91.3-95.1)	82.6 (79.7-85.6)
45-64	86.5 (85.0-88.1)	75.8 (73.8-77.9)	52.1 (49.7-54.5)
65-74	77.9 (75.7-80.1)	61.8 (59.2-64.4)	27.1 (24.7-29.7)
75+	69.7 (66.4-73.3)	45.9 (42.2-49.9)	14.1 (11.5-17.2)
Men	86.1 (84.9-87.3)	74.4 (72.8-76.0)	46.4 (44.5-48.5)
Women	87.7 (86.5-89.0)	76.9 (75.3-78.7)	50.7 (48.4-53.1)
Diabetes	81.5 (79.6-83.4)	64.9 (62.5-67.4)	31.5 (28.9-34.2)
Hypertension / renal vascular disease	87.4 (85.3-89.6)	74.3 (71.4-77.4)	46.2 (42.5-50.3)
Glomerulonephritis	89.9 (87.3-92.5)	81.4 (78.0-84.8)	58.3 (53.9-62.9)
Other cause	85.7 (84.2-87.2)	76.0 (74.2-77.9)	50.8 (48.4-53.2)
All	86.8 (85.9-87.6)	75.4 (74.3-76.6)	48.2 (46.6-49.8)

Table A.5.5

90-day, one- and two-year survival probabilities (cohort 1998-2002): Incident RRT patients, unadjusted
from day 1 to 90 days, one and two years, by age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES			
0-19	99.5 (98.9-99.8)	97.3 (96.1-98.1)	96.1 (94.8-97.1)
20-44	98.6 (98.4-98.8)	95.8 (95.5-96.1)	92.7 (92.2-93.1)
45-64	96.1 (95.9-96.3)	88.5 (88.2-88.8)	80.2 (79.8-80.6)
65-74	92.3 (92.0-92.6)	78.3 (77.9-78.7)	64.5 (64.1-64.9)
75+	88.2 (87.8-88.5)	68.8 (68.4-69.2)	51.1 (50.7-51.5)
Men	93.6 (93.4-93.8)	82.2 (81.9-82.5)	70.9 (70.7-71.2)
Women	93.4 (93.1-93.6)	81.7 (81.3-82.0)	70.2 (69.9-70.6)
Diabetes	94.1 (93.8-94.4)	80.3 (79.8-80.8)	64.9 (64.4-65.4)
Hypertension / renal vascular disease	92.5 (92.1-92.9)	79.0 (78.5-79.5)	65.5 (65.0-66.0)
Glomerulonephritis	97.0 (96.7-97.3)	91.5 (91.0-91.9)	85.4 (84.9-86.0)
Other cause	92.6 (92.4-92.8)	81.1 (80.8-81.4)	70.8 (70.5-71.1)
All	93.5 (93.3-93.6)	82.0 (81.8-82.2)	70.7 (70.4-70.9)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA			
0-19	100	98.6 (90.8-99.8)	97.2 (89.7-99.3)
20-44	97.6 (96.2-98.5)	94.1 (92.2-95.6)	91.4 (89.2-93.1)
45-64	94.7 (93.6-95.5)	84.3 (82.9-85.6)	73.7 (72.2-75.1)
65-74	89.0 (87.5-90.4)	72.0 (70.3-73.7)	54.5 (53.1-55.9)
75+	84.5 (82.5-86.3)	63.4 (61.4-65.2)	44.3 (43.0-45.7)
Men	92.7 (91.8-93.5)	79.8 (78.6-80.9)	67.5 (66.4-68.6)
Women	89.9 (88.6-91.0)	76.3 (74.9-77.7)	62.4 (61.0-63.7)
Diabetes	90.8 (89.5-92.0)	73.7 (72.1-75.2)	56.6 (55.2-57.9)
Hypertension / renal vascular disease	91.7 (89.8-93.2)	78.7 (76.4-80.8)	61.1 (59.0-63.2)
Glomerulonephritis	96.4 (94.8-97.5)	88.7 (86.4-90.6)	83.1 (80.6-85.2)
Other cause	90.5 (89.3-91.6)	78.6 (77.1-80.0)	68.7 (67.2-70.1)
All	91.6 (90.9-92.3)	78.4 (77.5-79.3)	65.5 (64.6-66.3)

Table A.5.6

90-day, one- and two-year survival probabilities (cohort 1998-2002): Incident RRT patients, adjusted from day 1 to 90 days, one and two years, adjusted for age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES			
0-19	99.5 (99.0-99.9)	97.0 (96.0-98.1)	95.8 (94.5-97.1)
20-44	98.6 (98.4-98.8)	95.7 (95.3-96.1)	92.5 (92.0-93.0)
45-64	96.3 (96.0-96.5)	88.9 (88.5-89.3)	80.8 (80.3-81.2)
65-74	92.6 (92.3-93.0)	79.3 (78.8-79.8)	66.0 (65.4-66.6)
75+	88.3 (87.9-88.8)	69.3 (68.7-70.0)	51.8 (51.1-52.5)
Men	95.6 (95.4-95.7)	87.3 (87.0-87.5)	78.2 (77.8-78.5)
Women	95.8 (95.6-95.9)	87.7 (87.4-88.0)	78.9 (78.5-79.3)
Diabetes	95.6 (95.3-95.9)	84.8 (84.3-85.3)	71.7 (71.1-72.3)
Hypertension / renal vascular disease	96.0 (95.7-96.2)	88.1 (87.6-88.5)	79.2 (78.6-79.8)
Glomerulonephritis	97.3 (97.0-97.6)	92.0 (91.5-92.5)	86.0 (85.4-86.6)
Other cause	94.9 (94.7-95.1)	86.3 (86.0-86.6)	77.8 (77.4-78.2)
All	95.7 (95.5-95.8)	87.4 (87.2-87.7)	78.5 (78.2-78.8)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	97.5 (96.3-98.7)	93.9 (92.1-95.7)	91.0 (88.8-93.2)
45-64	95.1 (94.2-96.0)	85.4 (83.9-86.9)	75.4 (73.5-77.3)
65-74	90.2 (88.7-91.6)	74.6 (72.4-76.7)	58.2 (55.8-60.7)
75+	85.4 (83.3-87.5)	65.7 (62.9-68.7)	47.1 (44.2-50.3)
Men	94.7 (94.0-95.4)	84.4 (83.3-85.5)	73.7 (72.2-75.1)
Women	94.1 (93.3-94.9)	85.4 (84.2-86.6)	74.9 (73.4-76.5)
Diabetes	93.5 (92.5-94.4)	80.2 (78.6-81.9)	65.6 (63.6-67.7)
Hypertension / renal vascular disease	95.1 (94.0-96.2)	87.1 (85.3-88.9)	74.5 (72.1-77.0)
Glomerulonephritis	96.1 (94.7-97.6)	87.5 (85.0-90.1)	81.0 (78.0-84.1)
Other cause	92.8 (91.8-93.8)	82.9 (81.4-84.5)	73.8 (72.0-75.7)
All	94.4 (93.9-94.9)	84.8 (84.0-85.6)	74.2 (73.1-75.2)

Table A.5.7

One- and two-year survival probabilities (cohort 1998-2002): Incident RRT patients, unadjusted
from day 91 to one and two years + 90 days, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	97.2 (96.0-98.1)	96.3 (94.9-97.3)
20-44	96.3 (96.0-96.6)	93.0 (92.5-93.4)
45-64	89.7 (89.4-90.0)	81.4 (81.0-81.8)
65-74	80.9 (80.5-81.3)	66.5 (66.1-66.9)
75+	72.6 (72.1-73.0)	53.6 (53.2-54.0)
Men	84.6 (84.3-84.9)	72.9 (72.6-73.2)
Women	83.9 (83.6-84.3)	72.4 (72.0-72.7)
Diabetes	80.8 (80.3-81.3)	65.1 (64.7-65.6)
Hypertension / renal vascular disease	81.6 (81.1-82.1)	67.5 (67.0-68.1)
Glomerulonephritis	92.6 (92.2-93.1)	86.3 (85.7-86.8)
Other cause	84.4 (84.1-84.7)	73.8 (73.5-74.1)
All	84.3 (84.1-84.6)	72.7 (72.5-72.9)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19	98.0 (87.2-99.7)	98.0 (87.2-99.7)
20-44	95.2 (93.3-96.6)	92.1 (89.8-93.9)
45-64	85.9 (84.4-87.2)	75.1 (73.5-76.6)
65-74	75.3 (73.5-77.1)	57.7 (56.1-59.3)
75+	68.9 (66.6-71.0)	46.7 (45.1-48.4)
Men	81.8 (80.5-82.9)	68.7 (67.5-69.9)
Women	80.5 (78.9-81.9)	65.9 (64.4-67.4)
Diabetes	76.0 (74.3-77.6)	57.9 (56.4-59.4)
Hypertension / renal vascular disease	81.9 (79.5-84.0)	63.0 (60.8-65.2)
Glomerulonephritis	90.3 (88.0-92.2)	84.0 (81.4-86.2)
Other cause	82.3 (80.7-83.7)	72.2 (70.6-73.7)
All	81.3 (80.3-82.2)	67.6 (66.7-68.6)

Table A.5.8

One- and two-year survival probabilities (cohort 1998-2002): Incident RRT patients, adjusted from day 91 to one and two years + 90 days, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	96.9 (95.8-98.1)	95.9 (94.5-97.2)
20-44	96.2 (95.9-96.6)	92.8 (92.4-93.3)
45-64	90.1 (89.8-90.5)	82.2 (81.7-82.6)
65-74	81.9 (81.4-82.4)	68.2 (67.6-68.8)
75+	73.0 (72.4-73.7)	54.5 (53.8-55.3)
Men	88.9 (88.6-89.2)	79.6 (79.3-80.0)
Women	89.1 (88.8-89.4)	80.4 (79.9-80.8)
Diabetes	85.1 (84.6-85.6)	71.9 (71.3-72.6)
Hypertension / renal vascular disease	89.5 (89.1-89.9)	80.4 (79.8-81.0)
Glomerulonephritis	93.0 (92.5-93.5)	86.7 (86.1-87.4)
Other cause	88.5 (88.2-88.8)	80.0 (79.6-80.4)
All	89.0 (88.8-89.2)	79.9 (79.6-80.2)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	95.0 (93.2-96.8)	91.8 (89.6-94.1)
45-64	87.1 (85.6-88.6)	77.3 (75.4-79.2)
65-74	78.1 (76.0-80.3)	62.0 (59.5-64.7)
75+	71.2 (68.3-74.3)	50.2 (46.9-53.7)
Men	86.2 (85.1-87.3)	75.3 (73.8-76.8)
Women	88.1 (86.9-89.3)	77.8 (76.2-79.4)
Diabetes	82.0 (80.3-83.7)	66.9 (64.8-69.1)
Hypertension / renal vascular disease	89.0 (87.2-90.7)	75.9 (73.3-78.5)
Glomerulonephritis	89.4 (86.9-91.9)	82.1 (79.0-85.3)
Other cause	85.7 (84.2-87.3)	77.0 (75.2-79.0)
All	87.0 (86.2-87.8)	76.3 (75.2-77.4)

Table A.5.9

90-day, one-, two- and five-year survival probabilities (cohort 1995-1999): Incident dialysis patients, unadjusted
from day 1 to 90 days, one, two and five years, by age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES				
0-19	99.0 (98.1-99.5)	96.5 (95.0-97.6)	94.8 (92.9-96.2)	87.9 (84.4-90.6)
20-44	98.7 (98.4-98.9)	95.2 (94.8-95.6)	91.1 (90.5-91.6)	78.3 (77.4-79.2)
45-64	96.4 (96.1-96.6)	88.3 (87.9-88.7)	78.8 (78.4-79.2)	53.6 (53.2-54.0)
65-74	92.3 (91.9-92.6)	78.3 (77.9-78.8)	63.8 (63.4-64.2)	32.2 (32.0-32.4)
75+	88.4 (87.9-88.9)	69.5 (68.9-70.0)	51.8 (51.3-52.2)	18.6 (18.5-18.8)
Men	94.0 (93.8-94.2)	82.7 (82.4-83.0)	70.3 (70.0-70.6)	40.5 (40.3-40.7)
Women	93.8 (93.5-94.0)	82.3 (81.9-82.7)	69.9 (69.5-70.3)	41.9 (41.6-42.1)
Diabetes	94.1 (93.7-94.5)	79.8 (79.2-80.3)	61.9 (61.4-62.4)	26.9 (26.7-27.1)
Hypertension / renal vascular disease	92.6 (92.1-93.0)	79.7 (79.1-80.2)	65.5 (64.9-66.1)	33.8 (33.5-34.1)
Glomerulonephritis	96.7 (96.3-97.0)	90.6 (90.1-91.1)	83.0 (82.4-83.7)	58.9 (58.2-59.6)
Other cause	93.5 (93.2-93.7)	82.2 (81.8-82.5)	71.4 (71.0-71.7)	44.9 (44.6-45.2)
All	93.9 (93.8-94.1)	82.5 (82.3-82.8)	70.1 (69.9-70.4)	41.0 (40.9-41.2)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA				
0-19	100	100	100	88.2 (65.3-96.4)
20-44	98.1 (96.8-98.9)	94.9 (93.0-96.3)	92.1 (89.8-93.9)	75.4 (71.7-78.7)
45-64	94.6 (93.5-95.5)	83.0 (81.5-84.4)	71.1 (69.5-72.7)	41.1 (40.0-42.2)
65-74	90.0 (88.5-91.4)	71.5 (69.7-73.1)	54.9 (53.4-56.4)	20.5 (20.0-21.0)
75+	86.5 (84.3-88.5)	63.7 (61.4-65.8)	42.4 (40.8-43.9)	11.1 (10.9-11.4)
Men	93.4 (92.4-94.2)	79.5 (78.2-80.6)	65.6 (64.3-66.8)	32.7 (32.0-33.4)
Women	91.3 (90.1-92.4)	76.5 (75.0-77.9)	62.5 (61.0-63.9)	31.4 (30.7-32.2)
Diabetes	91.3 (89.8-92.5)	73.5 (71.7-75.1)	53.8 (52.3-55.2)	18.7 (18.3-19.1)
Hypertension / renal vascular disease	93.4 (91.3-94.9)	78.4 (75.7-80.8)	60.8 (58.4-63.1)	27.8 (26.7-28.8)
Glomerulonephritis	96.3 (94.6-97.4)	88.9 (86.5-90.9)	82.6 (79.9-84.9)	57.5 (54.7-60.1)
Other cause	91.8 (90.7-92.9)	78.1 (76.5-79.5)	67.4 (65.9-68.8)	37.4 (36.5-38.4)
All	92.5 (91.8-93.2)	78.2 (77.3-79.2)	64.3 (63.3-65.2)	32.2 (31.7-32.7)

Table A.5.10
90-day, one-, two- and five-year survival probabilities (cohort 1995-1999): Incident dialysis patients, adjusted from day 1 to 90 days, one, two and five years, adjusted for age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES				
0-19	98.9 (98.2-99.6)	96.1 (94.7-97.6)	94.1 (92.1-96.1)	84.9 (80.5-89.6)
20-44	98.6 (98.4-98.9)	95.2 (94.8-95.7)	90.9 (90.3-91.6)	77.0 (75.8-78.2)
45-64	96.5 (96.3-96.8)	88.9 (88.5-89.3)	79.7 (79.2-80.3)	54.1 (53.4-54.9)
65-74	92.6 (92.3-93.0)	79.5 (78.9-80.0)	65.4 (64.7-66.0)	33.5 (32.8-34.2)
75+	88.5 (88.0-89.1)	69.9 (69.1-70.7)	52.4 (51.5-53.3)	18.8 (18.1-19.5)
Men	95.4 (95.2-95.5)	86.4 (86.0-86.7)	76.1 (75.7-76.5)	48.2 (47.6-48.8)
Women	95.4 (95.2-95.7)	86.8 (86.4-87.2)	77.0 (76.5-77.5)	51.4 (50.8-52.1)
Diabetes	95.1 (94.7-95.4)	82.8 (82.2-83.4)	66.9 (66.1-67.7)	31.9 (31.0-32.8)
Hypertension / renal vascular disease	95.3 (95.0-95.6)	86.8 (86.2-87.3)	76.6 (75.9-77.3)	48.9 (47.9-49.9)
Glomerulonephritis	96.7 (96.4-97.1)	90.9 (90.3-91.4)	83.6 (82.8-84.3)	59.3 (58.1-60.5)
Other cause	94.8 (94.6-95.0)	85.7 (85.3-86.0)	76.6 (76.2-77.1)	52.2 (51.6-52.8)
All	95.4 (95.3-95.5)	86.5 (86.3-86.8)	76.5 (76.1-76.8)	49.5 (49.0-50.0)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA				
0-19				
20-44	98.1 (97.1-99.1)	94.9 (93.2-96.6)	91.9 (89.7-94.2)	71.5 (65.9-77.5)
45-64	95.0 (94.0-96.0)	84.3 (82.7-86.0)	73.1 (71.1-75.3)	42.1 (39.3-45.0)
65-74	91.0 (89.6-92.4)	74.6 (72.5-76.8)	59.2 (56.7-61.8)	23.6 (21.3-26.1)
75+	87.6 (85.3-89.9)	66.3 (63.1-69.7)	45.5 (42.1-49.2)	13.3 (11.0-16.1)
Men	94.7 (94.0-95.5)	83.5 (82.2-84.7)	71.3 (69.6-72.9)	38.1 (35.9-40.3)
Women	94.1 (93.3-95.0)	83.6 (82.2-85.0)	72.7 (70.9-74.5)	42.8 (40.3-45.4)
Diabetes	93.1 (91.9-94.3)	78.6 (76.7-80.6)	61.2 (58.8-63.7)	25.2 (22.7-27.9)
Hypertension / renal vascular disease	95.6 (94.3-96.9)	85.0 (82.8-87.4)	71.2 (68.1-74.4)	39.1 (35.2-43.4)
Glomerulonephritis	95.6 (94.0-97.3)	86.9 (84.2-89.8)	79.6 (76.2-83.2)	49.4 (44.1-55.3)
Other cause	93.1 (92.1-94.2)	81.3 (79.7-83.0)	71.8 (69.8-73.8)	41.6 (39.0-44.3)
All	94.4 (93.9-95.0)	83.5 (82.5-84.4)	71.8 (70.6-73.0)	40.0 (38.3-41.7)

Table A.5.11
One-, two- and five-year survival probabilities (cohort 1995-1999): Incident dialysis patients, unadjusted
from day 91 to one, two and five years + 90 days, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	97.4 (95.9-98.3)	95.5 (93.4-96.9)	87.7 (83.9-90.6)
20-44	95.6 (95.1-96.0)	91.2 (90.6-91.8)	77.9 (77.0-78.8)
45-64	89.0 (88.6-89.4)	79.5 (79.0-79.9)	53.3 (52.9-53.7)
65-74	80.9 (80.4-81.3)	65.5 (65.1-66.0)	32.6 (32.4-32.8)
75+	73.4 (72.8-74.0)	54.2 (53.7-54.7)	19.1 (18.9-19.2)
Men	84.6 (84.3-85.0)	71.6 (71.3-72.0)	40.7 (40.4-40.9)
Women	84.3 (83.9-84.7)	71.6 (71.1-72.0)	42.3 (42.0-42.6)
Diabetes	79.9 (79.3-80.5)	61.8 (61.3-62.4)	26.1 (25.9-26.3)
Hypertension / renal vascular disease	82.1 (81.5-82.7)	67.2 (66.6-67.8)	34.2 (33.9-34.5)
Glomerulonephritis	91.9 (91.4-92.4)	83.9 (83.2-84.5)	58.5 (57.8-59.2)
Other cause	85.0 (84.6-85.3)	73.5 (73.2-73.9)	45.8 (45.5-46.0)
All	84.5 (84.3-84.8)	71.6 (71.3-71.9)	41.3 (41.1-41.5)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19	100	100	87.1 (63.0-96.0)
20-44	95.8 (93.9-97.1)	92.7 (90.3-94.5)	74.5 (70.6-78.0)
45-64	84.8 (83.2-86.2)	72.3 (70.6-73.9)	41.6 (40.4-42.8)
65-74	74.5 (72.6-76.3)	56.6 (55.0-58.2)	21.0 (20.5-21.5)
75+	66.9 (64.4-69.3)	42.1 (40.5-43.8)	11.1 (10.8-11.4)
Men	81.2 (79.8-82.4)	66.2 (64.9-67.5)	33.3 (32.5-34.0)
Women	80.1 (78.5-81.5)	64.5 (62.9-66.0)	32.0 (31.1-32.8)
Diabetes	75.0 (73.1-76.8)	54.3 (52.7-55.8)	18.2 (17.8-18.6)
Hypertension / renal vascular disease	79.5 (76.7-81.9)	60.9 (58.4-63.3)	28.1 (27.0-29.2)
Glomerulonephritis	91.2 (88.8-93.0)	83.4 (80.6-85.9)	57.4 (54.4-60.2)
Other cause	81.6 (80.0-83.1)	69.6 (67.9-71.1)	39.1 (38.0-40.2)
All	80.7 (79.7-81.7)	65.5 (64.5-66.5)	32.7 (32.2-33.3)

Table A.5.12

One-, two- and five-year survival probabilities (cohort 1995-1999): Incident dialysis patients, adjusted from day 91 to one, two and five years + 90 days, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	97.0 (95.7-98.4)	94.8 (92.8-96.9)	84.7 (80.0-89.8)
20-44	95.5 (95.1-96.0)	91.0 (90.3-91.7)	76.5 (75.2-77.8)
45-64	89.5 (89.1-89.9)	80.1 (79.6-80.7)	53.7 (52.9-54.5)
65-74	81.7 (81.1-82.2)	66.7 (66.0-67.3)	33.6 (32.9-34.3)
75+	73.4 (72.6-74.2)	54.1 (53.2-55.1)	19.0 (18.2-19.7)
Men	87.7 (87.4-88.0)	76.8 (76.3-77.2)	47.9 (47.3-48.5)
Women	88.0 (87.7-88.4)	77.8 (77.3-78.3)	51.2 (50.5-52.0)
Diabetes	82.8 (82.1-83.4)	66.6 (65.8-67.5)	31.2 (30.3-32.2)
Hypertension / renal vascular disease	88.2 (87.7-88.7)	77.4 (76.7-78.2)	49.2 (48.2-50.2)
Glomerulonephritis	92.1 (91.5-92.6)	84.4 (83.6-85.2)	59.3 (58.1-60.5)
Other cause	87.8 (87.5-88.2)	78.2 (77.8-78.7)	53.0 (52.4-53.7)
All	87.8 (87.6-88.1)	77.2 (76.8-77.5)	49.2 (48.7-49.7)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	95.7 (94.0-97.4)	92.3 (90.0-94.7)	69.4 (63.2-76.1)
45-64	86.0 (84.4-87.7)	74.1 (72.0-76.3)	42.4 (39.6-45.5)
65-74	77.3 (75.1-79.6)	60.7 (58.1-63.4)	24.0 (21.5-26.7)
75+	69.2 (65.8-72.8)	45.1 (41.4-49.1)	13.5 (11.0-16.6)
Men	84.6 (83.3-85.9)	71.5 (69.7-73.2)	38.2 (35.9-40.6)
Women	86.0 (84.6-87.4)	73.7 (71.9-75.7)	42.6 (39.9-45.4)
Diabetes	79.7 (77.7-81.8)	61.6 (59.0-64.2)	24.4 (21.8-27.3)
Hypertension / renal vascular disease	85.7 (83.3-88.1)	71.0 (67.8-74.4)	39.2 (35.1-43.7)
Glomerulonephritis	89.6 (87.0-92.3)	80.7 (77.2-84.4)	50.1 (44.6-56.3)
Other cause	84.2 (82.6-85.9)	73.2 (71.2-75.4)	42.7 (40.0-45.7)
All	85.1 (84.2-86.1)	72.4 (71.1-73.7)	39.9 (38.2-41.8)

Table A.5.13
90-day, one- and two-year survival probabilities (cohort 1998-2002): Incident dialysis patients, unadjusted
from day 1 to 90 days, one and two years, by age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES			
0-19	99.4 (98.6-99.8)	97.2 (95.8-98.2)	95.9 (94.0-97.2)
20-44	98.6 (98.3-98.8)	95.4 (95.0-95.8)	91.3 (90.8-91.8)
45-64	96.1 (95.9-96.3)	88.2 (87.8-88.5)	78.9 (78.5-79.3)
65-74	92.2 (91.9-92.5)	78.1 (77.7-78.5)	64.2 (63.8-64.5)
75+	88.0 (87.6-88.4)	68.7 (68.2-69.1)	51.0 (50.6-51.3)
Men	93.4 (93.2-93.6)	81.6 (81.3-81.8)	69.2 (68.9-69.4)
Women	93.2 (93.0-93.5)	81.0 (80.7-81.4)	68.5 (68.2-68.9)
Diabetes	94.0 (93.7-94.3)	79.9 (79.4-80.4)	63.5 (63.1-64.0)
Hypertension / renal vascular disease	92.5 (92.1-92.8)	78.8 (78.3-79.3)	64.8 (64.3-65.3)
Glomerulonephritis	96.9 (96.6-97.2)	91.0 (90.5-91.4)	83.8 (83.1-84.3)
Other cause	92.4 (92.2-92.6)	80.4 (80.1-80.7)	68.9 (68.6-69.2)
All	93.3 (93.2-93.5)	81.4 (81.2-81.6)	68.9 (68.7-69.1)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA			
0-19	100	100	100
20-44	97.7 (96.2-98.6)	93.4 (91.3-95.0)	90.5 (88.0-92.5)
45-64	94.6 (93.5-95.4)	84.0 (82.5-85.3)	72.2 (70.6-73.7)
65-74	89.1 (87.6-90.4)	71.9 (70.1-73.5)	53.8 (52.4-55.2)
75+	84.5 (82.5-86.3)	63.4 (61.4-65.2)	44.3 (42.9-45.7)
Men	92.6 (91.7-93.4)	78.9 (77.7-80.1)	65.2 (64.0-66.4)
Women	89.7 (88.5-90.8)	75.7 (74.2-77.1)	60.5 (59.1-61.8)
Diabetes	90.6 (89.3-91.8)	73.1 (71.4-74.6)	54.4 (53.0-55.7)
Hypertension / renal vascular disease	91.6 (89.7-93.2)	78.5 (76.1-80.6)	60.4 (58.3-62.5)
Glomerulonephritis	96.2 (94.5-97.4)	87.8 (85.4-89.8)	81.5 (78.8-83.9)
Other cause	90.3 (89.1-91.5)	77.7 (76.2-79.2)	66.5 (64.9-67.9)
All	91.4 (90.7-92.1)	77.6 (76.7-78.5)	63.3 (62.4-64.2)

Table A.5.14
90-day, one- and two-year survival probabilities (cohort 1998-2002): Incident dialysis patients, adjusted from day 1 to 90 days, one and two years, adjusted for age, gender and primary diagnosis

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES			
0-19	99.4 (98.8-99.9)	97.0 (95.6-98.3)	95.3 (93.5-97.2)
20-44	98.5 (98.3-98.8)	95.3 (94.9-95.7)	91.1 (90.5-91.7)
45-64	96.2 (96.0-96.4)	88.6 (88.2-89.0)	79.6 (79.1-80.1)
65-74	92.5 (92.2-92.8)	79.1 (78.6-79.6)	65.5 (64.9-66.1)
75+	88.1 (87.7-88.6)	69.1 (68.5-69.8)	51.5 (50.8-52.3)
Men	95.2 (95.1-95.4)	86.2 (86.0-86.5)	76.3 (75.9-76.7)
Women	95.3 (95.1-95.5)	86.6 (86.3-86.9)	77.0 (76.5-77.5)
Diabetes	95.3 (95.0-95.6)	83.9 (83.4-84.4)	69.9 (69.3-70.6)
Hypertension / renal vascular disease	95.5 (95.3-95.8)	86.9 (86.5-87.4)	77.3 (76.7-77.9)
Glomerulonephritis	97.1 (96.8-97.4)	91.5 (91.0-92.0)	84.8 (84.1-85.5)
Other cause	94.4 (94.2-94.6)	85.2 (84.9-85.5)	75.9 (75.5-76.3)
All	95.3 (95.1-95.4)	86.4 (86.2-86.6)	76.6 (76.3-76.9)

	90 day Survival (95% CI)	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	97.5 (96.3-98.8)	93.1 (91.1-95.2)	90.0 (87.4-92.7)
45-64	94.9 (94.0-95.9)	85.0 (83.5-86.6)	73.9 (71.9-75.9)
65-74	90.1 (88.6-91.5)	74.1 (72.0-76.4)	57.1 (54.6-59.7)
75+	85.2 (83.1-87.4)	65.4 (62.6-68.4)	46.7 (43.7-49.9)
Men	94.3 (93.6-95.0)	83.3 (82.1-84.5)	71.6 (70.1-73.2)
Women	93.5 (92.7-94.4)	84.0 (82.7-85.3)	72.5 (70.8-74.2)
Diabetes	93.0 (91.9-94.0)	79.1 (77.3-80.8)	63.3 (61.1-65.5)
Hypertension / renal vascular disease	94.6 (93.4-95.9)	85.9 (84.0-87.9)	72.5 (69.9-75.2)
Glomerulonephritis	95.9 (94.4-97.5)	86.8 (84.2-89.6)	80.1 (76.8-83.4)
Other cause	92.3 (91.3-93.4)	81.7 (80.1-83.4)	71.7 (69.8-73.7)
All	93.9 (93.4-94.5)	83.6 (82.6-84.5)	71.9 (70.7-73.1)

Table A.5.15
One- and two-year survival probabilities (cohort 1998-2002): Incident dialysis patients, unadjusted
from day 91 to one and two years + 90 days, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	97.7 (96.3-98.6)	96.2 (94.2-97.5)
20-44	95.9 (95.5-96.2)	91.4 (90.8-91.9)
45-64	89.3 (88.9-89.6)	79.9 (79.5-80.3)
65-74	80.7 (80.3-81.2)	66.1 (65.6-66.5)
75+	72.6 (72.1-73.0)	53.5 (53.1-53.9)
Men	83.9 (83.7-84.2)	70.8 (70.5-71.1)
Women	83.3 (83.0-83.7)	70.5 (70.1-70.9)
Diabetes	80.4 (79.9-80.9)	63.6 (63.1-64.1)
Hypertension / renal vascular disease	81.4 (80.9-81.9)	66.6 (66.1-67.2)
Glomerulonephritis	92.1 (91.6-92.6)	84.3 (83.7-85.0)
Other cause	83.7 (83.3-84.0)	71.7 (71.3-72.0)
All	83.7 (83.5-83.9)	70.7 (70.5-70.9)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19	100	100
20-44	95.0 (92.9-96.5)	91.2 (88.5-93.3)
45-64	85.6 (84.1-87.0)	73.9 (72.2-75.4)
65-74	75.0 (73.2-76.8)	56.7 (55.1-58.3)
75+	68.9 (66.6-71.0)	46.7 (45.0-48.3)
Men	81.3 (80.0-82.5)	66.7 (65.5-68.0)
Women	80.0 (78.4-81.4)	64.2 (62.7-65.7)
Diabetes	75.5 (73.7-77.1)	55.7 (54.2-57.2)
Hypertension / renal vascular disease	81.7 (79.3-83.9)	62.1 (59.8-64.3)
Glomerulonephritis	89.9 (87.5-91.9)	83.0 (80.2-85.4)
Other cause	81.7 (80.0-83.2)	70.5 (68.8-72.1)
All	80.8 (79.8-81.7)	65.7 (64.7-66.7)

Table A.5.16

One- and two-year survival probabilities (cohort 1998-2002): Incident dialysis patients, adjusted from day 91 to one and two years + 90 days, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	97.5 (96.3-98.8)	95.8 (94.0-97.7)
20-44	95.8 (95.4-96.2)	91.2 (90.6-91.8)
45-64	89.7 (89.4-90.1)	80.8 (80.3-81.3)
65-74	81.7 (81.2-82.2)	67.7 (67.1-68.3)
75+	72.9 (72.2-73.6)	54.2 (53.5-55.0)
Men	87.9 (87.6-88.2)	77.6 (77.2-78.0)
Women	88.1 (87.8-88.4)	78.4 (77.9-78.9)
Diabetes	84.1 (83.6-84.6)	70.0 (69.3-70.7)
Hypertension / renal vascular disease	88.4 (87.9-88.8)	78.4 (77.7-79.0)
Glomerulonephritis	92.6 (92.1-93.1)	85.4 (84.7-86.1)
Other cause	87.5 (87.2-87.8)	78.0 (77.5-78.4)
All	88.0 (87.8-88.2)	78.0 (77.6-78.3)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	94.8 (92.9-96.7)	90.7 (88.1-93.5)
45-64	86.7 (85.2-88.3)	76.1 (74.1-78.1)
65-74	77.7 (75.5-79.9)	60.8 (58.2-63.6)
75+	70.9 (67.9-74.0)	49.6 (46.3-53.1)
Men	85.2 (84.0-86.4)	73.1 (71.5-74.7)
Women	86.9 (85.6-88.2)	75.5 (73.7-77.3)
Diabetes	80.9 (79.1-82.7)	64.5 (62.2-66.9)
Hypertension / renal vascular disease	87.9 (86.0-89.9)	73.7 (71.0-76.5)
Glomerulonephritis	89.1 (86.6-91.8)	81.5 (78.2-85.0)
Other cause	84.6 (82.9-86.2)	75.0 (72.9-77.0)
All	85.9 (85.0-86.8)	74.1 (72.9-75.3)

A5 SURVIVAL PROBABILITY
Patient Survival after First Transplant (cadaver donor)

Table A.5.17
One-, two- and five-year survival probabilities (cohort 1995-1999): First transplant patients (cadaver donor), unadjusted from day of transplant to one, two and five years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	97.3 (95.6-98.3)	96.2 (94.4-97.5)	94.4 (92.3-96.0)
20-44	97.7 (97.2-98.1)	96.6 (96.1-97.1)	93.0 (92.3-93.7)
45-64	94.2 (93.6-94.8)	91.6 (90.9-92.2)	82.3 (81.4-83.1)
65+	86.1 (83.5-88.3)	81.5 (78.8-83.9)	62.8 (60.3-65.2)
Men	95.2 (94.7-95.7)	93.1 (92.5-93.6)	85.6 (84.8-86.3)
Women	95.5 (94.9-96.1)	93.7 (93.0-94.4)	87.4 (86.5-88.3)
Diabetes	92.5 (91.0-93.8)	90.1 (88.4-91.5)	79.3 (77.4-81.1)
Hypertension / renal vascular disease	93.3 (91.6-94.7)	89.6 (87.7-91.3)	79.1 (76.9-81.2)
Glomerulonephritis	96.8 (96.1-97.3)	95.2 (94.4-95.9)	89.5 (88.5-90.4)
Other cause	95.4 (94.9-95.9)	93.6 (92.9-94.1)	87.0 (86.2-87.8)
All	95.3 (95.0-95.7)	93.3 (92.9-93.8)	86.3 (85.7-86.8)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19	96.6 (87.7-99.1)	93.2 (83.9-97.2)	89.8 (80.2-94.9)
20-44	97.4 (95.6-98.4)	95.4 (93.2-96.8)	92.7 (90.3-94.6)
45-64	91.9 (89.5-93.7)	87.4 (84.8-89.7)	77.8 (74.9-80.4)
65+	82.1 (73.8-87.9)	75.6 (67.5-82.0)	65.4 (58.0-71.8)
Men	94.0 (92.1-95.4)	90.4 (88.2-92.1)	83.6 (81.2-85.7)
Women	93.4 (91.0-95.3)	90.2 (87.4-92.4)	84.3 (81.2-86.9)
Diabetes	94.5 (90.0-97.0)	89.6 (84.5-93.1)	78.5 (73.0-83.0)
Hypertension / renal vascular disease	88.0 (80.1-92.8)	80.7 (72.7-86.6)	72.3 (64.6-78.6)
Glomerulonephritis	96.8 (94.5-98.2)	94.6 (91.8-96.4)	90.3 (87.0-92.7)
Other cause	92.6 (90.3-94.3)	89.3 (86.8-91.3)	83.2 (80.5-85.6)
All	93.8 (92.3-95.0)	90.3 (88.6-91.7)	83.9 (82.0-85.6)

A5 SURVIVAL PROBABILITY
Patient Survival after First Transplant (cadaver donor)

Table A.5.18

One-, two- and five-year survival probabilities (cohort 1995-1999): First transplant patients (cadaver donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	97.1 (95.6-98.5)	95.9 (94.2-97.6)	93.9 (91.9-96.0)
20-44	97.8 (97.4-98.2)	96.8 (96.3-97.3)	93.4 (92.7-94.1)
45-64	94.4 (93.8-95.0)	91.9 (91.2-92.6)	82.9 (81.8-83.9)
65+	86.4 (83.6-89.1)	82.0 (79.0-85.2)	63.6 (59.8-67.6)
Men	96.4 (96.0-96.8)	94.7 (94.2-95.2)	88.5 (87.8-89.3)
Women	96.3 (95.7-96.8)	94.8 (94.2-95.5)	89.4 (88.5-90.3)
Diabetes	93.1 (91.7-94.5)	90.9 (89.3-92.5)	80.7 (78.5-83.0)
Hypertension / renal vascular disease	95.9 (94.9-96.9)	93.7 (92.4-95.0)	86.7 (84.9-88.6)
Glomerulonephritis	97.3 (96.8-97.8)	95.9 (95.3-96.6)	90.8 (89.9-91.8)
Other cause	96.3 (95.9-96.8)	94.8 (94.2-95.3)	89.2 (88.4-90.0)
All	96.4 (96.0-96.7)	94.8 (94.4-95.2)	88.9 (88.3-89.5)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	97.6 (96.3-98.9)	95.7 (94.0-97.4)	93.2 (91.1-95.4)
45-64	92.1 (89.9-94.4)	88.1 (85.4-90.8)	78.8 (75.5-82.3)
65+	83.4 (75.9-91.7)	77.4 (69.0-86.8)	67.8 (58.5-78.5)
Men	95.5 (94.2-96.9)	92.8 (91.1-94.4)	87.3 (85.2-89.6)
Women	94.5 (92.6-96.4)	91.9 (89.7-94.2)	86.8 (84.0-89.7)
Diabetes	94.7 (91.6-98.0)	91.0 (86.9-95.1)	80.9 (75.4-86.8)
Hypertension / renal vascular disease			
Glomerulonephritis	97.0 (95.3-98.8)	94.8 (92.5-97.1)	90.5 (87.5-93.6)
Other cause	94.4 (92.7-96.0)	91.8 (89.8-93.8)	86.9 (84.4-89.4)
All	95.2 (94.1-96.3)	92.5 (91.1-93.8)	87.2 (85.4-89.0)

Table A.5.19

One- and two-year survival probabilities (cohort 1998-2002): First transplant patients (cadaver donor), unadjusted
from day of transplant to one and two years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	98.0 (96.4-98.9)	97.0 (95.2-98.2)
20-44	97.8 (97.3-98.1)	96.7 (96.2-97.2)
45-64	95.1 (94.6-95.6)	92.8 (92.1-93.3)
65+	89.3 (87.3-90.9)	84.5 (82.4-86.4)
Men	95.7 (95.2-96.1)	93.7 (93.1-94.1)
Women	96.1 (95.6-96.6)	94.3 (93.6-94.9)
Diabetes	94.3 (93.1-95.3)	91.6 (90.2-92.7)
Hypertension / renal vascular disease	93.6 (92.1-94.9)	90.8 (89.1-92.2)
Glomerulonephritis	96.7 (96.1-97.3)	94.9 (94.2-95.6)
Other cause	96.1 (95.7-96.6)	94.4 (93.9-94.9)
All	95.8 (95.5-96.2)	93.9 (93.5-94.3)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19	97.6 (84.6-99.6)	92.7 (80.5-97.4)
20-44	97.0 (95.2-98.1)	94.8 (92.6-96.3)
45-64	92.6 (90.5-94.3)	89.6 (87.2-91.5)
65+	83.6 (77.2-88.3)	80.3 (73.9-85.3)
Men	93.1 (91.3-94.5)	90.3 (88.3-91.9)
Women	94.8 (92.5-96.4)	92.1 (89.5-94.0)
Diabetes	93.1 (89.4-95.5)	89.4 (85.4-92.4)
Hypertension / renal vascular disease	91.3 (85.2-95.0)	87.0 (80.5-91.4)
Glomerulonephritis	95.7 (93.2-97.3)	93.0 (90.1-95.0)
Other cause	93.2 (91.1-94.9)	91.1 (88.7-92.9)
All	93.7 (92.4-94.8)	90.9 (89.4-92.2)

Table A.5.20
One- and two-year survival probabilities (cohort 1998-2002): First transplant patients (cadaver donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19		
20-44	97.9 (97.5-98.3)	96.9 (96.4-97.4)
45-64	95.3 (94.8-95.8)	93.1 (92.4-93.7)
65+	89.6 (87.7-91.6)	85.2 (83.0-87.5)
Men	96.8 (96.4-97.2)	95.3 (94.8-95.7)
Women	97.0 (96.5-97.4)	95.5 (95.0-96.1)
Diabetes	95.1 (94.1-96.1)	92.7 (91.5-93.9)
Hypertension / renal vascular disease	96.1 (95.2-97.0)	94.4 (93.3-95.5)
Glomerulonephritis	97.3 (96.8-97.8)	95.8 (95.1-96.4)
Other cause	97.0 (96.6-97.4)	95.6 (95.1-96.1)
All	96.9 (96.6-97.2)	95.4 (95.0-95.7)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	97.2 (95.9-98.6)	95.2 (93.4-96.9)
45-64	93.0 (91.1-94.9)	90.4 (88.2-92.6)
65+	85.1 (79.2-91.3)	82.0 (75.8-88.8)
Men	95.1 (93.9-96.3)	93.0 (91.5-94.5)
Women	96.0 (94.6-97.6)	94.2 (92.4-96.0)
Diabetes	94.3 (91.8-96.9)	91.7 (88.7-94.8)
Hypertension / renal vascular disease		
Glomerulonephritis	96.3 (94.5-98.1)	93.9 (91.6-96.2)
Other cause	94.9 (93.3-96.4)	93.2 (91.4-95.0)
All	95.4 (94.5-96.4)	93.5 (92.3-94.6)

A5 SURVIVAL PROBABILITY
Patient Survival after First Transplant (living donor)

Table A.5.21
One-, two- and five-year survival probabilities (cohort 1995-1999): First transplant patients (living donor), unadjusted
from day of transplant to one, two and five years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	97.9 (95.5-99.0)	97.6 (95.1-98.8)	97.2 (94.6-98.6)
20-44	98.9 (98.2-99.4)	98.3 (97.5-98.9)	95.7 (94.4-96.6)
45-64	95.9 (94.1-97.1)	95.6 (93.8-96.8)	90.6 (88.4-92.5)
65+	90.2 (80.8-95.1)	86.9 (77.4-92.6)	78.7 (69.2-85.5)
Men	98.0 (97.2-98.6)	97.7 (96.8-98.4)	94.3 (93.1-95.4)
Women	97.0 (95.6-97.9)	95.9 (94.4-97.1)	93.1 (91.3-94.6)
Diabetes	97.8 (94.3-99.1)	96.7 (92.9-98.4)	86.0 (81.0-89.8)
Hypertension / renal vascular disease	91.1 (85.6-94.6)	89.6 (83.9-93.4)	85.9 (79.9-90.2)
Glomerulonephritis	98.6 (97.5-99.3)	98.2 (97.0-99.0)	95.8 (94.1-97.0)
Other cause	97.7 (96.7-98.4)	97.3 (96.2-98.1)	94.9 (93.5-96.0)
All	97.6 (96.9-98.2)	97.1 (96.3-97.7)	93.9 (92.9-94.8)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	98.5 (89.9-99.8)	98.5 (89.9-99.8)	96.9 (88.7-99.2)
45-64			
65+			
Men	97.1 (89.2-99.2)	97.1 (89.2-99.2)	95.6 (87.5-98.5)
Women	97.7 (85.2-99.7)	97.7 (85.2-99.7)	93.0 (81.3-97.5)
Diabetes			
Hypertension / renal vascular disease			
Glomerulonephritis	100	100	97.1 (82.1-99.6)
Other cause	96.9 (88.6-99.2)	96.9 (88.6-99.2)	93.7 (85.0-97.4)
All	97.3 (92.1-99.1)	97.3 (92.1-99.1)	94.6 (88.8-97.4)

A5 SURVIVAL PROBABILITY
Patient Survival after First Transplant (living donor)

Table A.5.22

One-, two- and five-year survival probabilities (cohort 1995-1999): First transplant patients (living donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19			
20-44	98.9 (98.4-99.5)	98.4 (97.7-99.1)	95.7 (94.6-96.9)
45-64	95.9 (94.4-97.4)	95.6 (94.1-97.2)	90.7 (88.5-93.0)
65+			
Men	97.9 (97.1-98.7)	97.6 (96.7-98.4)	93.8 (92.4-95.1)
Women	96.4 (95.0-97.8)	95.2 (93.6-96.8)	91.8 (89.8-94.0)
Diabetes	97.5 (95.2-99.9)	96.3 (93.4-99.2)	84.5 (79.1-90.3)
Hypertension / renal vascular disease			
Glomerulonephritis	98.4 (97.4-99.4)	97.9 (96.8-99.1)	95.0 (93.3-96.8)
Other cause	97.3 (96.3-98.3)	96.8 (95.7-97.9)	93.8 (92.3-95.4)
All	97.4 (96.6-98.1)	96.7 (95.9-97.5)	93.1 (91.9-94.2)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44			
45-64			
65+			
Men			
Women			
Diabetes			
Hypertension / renal vascular disease			
Glomerulonephritis			
Other cause			
All			

Table A.5.23

One- and two-year survival probabilities (cohort 1998-2002): First transplant patients (living donor), unadjusted
from day of transplant to one and two years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	97.9 (95.6-99.0)	97.9 (95.6-99.0)
20-44	98.9 (98.3-99.3)	98.5 (97.7-99.0)
45-64	96.3 (94.9-97.3)	95.2 (93.7-96.4)
65+	90.2 (83.1-94.5)	88.0 (80.7-92.7)
Men	97.4 (96.6-98.0)	96.8 (95.9-97.5)
Women	98.2 (97.2-98.9)	97.4 (96.2-98.2)
Diabetes	95.8 (92.3-97.7)	94.4 (90.7-96.7)
Hypertension / renal vascular disease	96.3 (92.6-98.2)	95.3 (91.4-97.4)
Glomerulonephritis	98.6 (97.6-99.2)	98.0 (96.9-98.8)
Other cause	97.6 (96.7-98.2)	97.0 (96.0-97.7)
All	97.7 (97.1-98.2)	97.0 (96.3-97.6)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	98.9 (92.7-99.8)	97.8 (91.8-99.4)
45-64	98.0 (87.1-99.7)	98.0 (87.1-99.7)
65+		
Men	98.4 (93.8-99.6)	97.6 (92.8-99.2)
Women	98.1 (87.6-99.7)	98.1 (87.6-99.7)
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis	98.4 (89.4-99.8)	96.8 (88.2-99.1)
Other cause	99.0 (93.1-99.9)	99.0 (93.1-99.9)
All	98.3 (94.9-99.4)	97.7 (94.2-99.1)

Table A.5.24
One- and two-year survival probabilities (cohort 1998-2002): First transplant patients (living donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19		
20-44	99.0 (98.5-99.5)	98.5 (97.9-99.1)
45-64	96.6 (95.4-97.7)	95.5 (94.2-96.9)
65+		
Men	97.3 (96.5-98.1)	96.7 (95.8-97.5)
Women	98.1 (97.2-99.0)	97.2 (96.1-98.3)
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis	98.5 (97.6-99.3)	97.9 (96.8-98.9)
Other cause	97.3 (96.4-98.2)	96.6 (95.6-97.6)
All	97.6 (97.0-98.2)	96.9 (96.2-97.6)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44		
45-64		
65+		
Men		
Women		
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis		
Other cause		
All		

A5 SURVIVAL PROBABILITY
Graft Survival after First Transplant (cadaver donor)

Table A.5.25
One-, two- and five-year survival probabilities (cohort 1995-1999): First graft (cadaver donor), unadjusted
from day of transplant to one, two and five years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	87.1 (84.5-89.3)	83.5 (80.7-85.9)	74.3 (71.5-76.9)
20-44	90.7 (89.9-91.4)	88.3 (87.4-89.1)	80.0 (79.0-80.9)
45-64	87.8 (87.0-88.6)	84.3 (83.5-85.1)	73.8 (72.9-74.7)
65+	81.1 (78.4-83.6)	75.1 (72.4-77.7)	57.5 (55.1-59.7)
Men	89.0 (88.3-89.7)	85.7 (85.0-86.4)	75.5 (74.7-76.3)
Women	87.9 (87.0-88.8)	84.9 (84.0-85.8)	75.4 (74.4-76.4)
Diabetes	86.6 (84.8-88.2)	83.0 (81.1-84.7)	70.2 (68.3-72.0)
Hypertension / renal vascular disease	86.1 (84.0-88.0)	81.0 (78.8-83.0)	69.3 (67.2-71.4)
Glomerulonephritis	90.3 (89.3-91.2)	87.6 (86.5-88.5)	77.7 (76.5-78.8)
Other cause	88.5 (87.7-89.2)	85.4 (84.6-86.1)	76.3 (75.4-77.1)
All	88.6 (88.1-89.1)	85.4 (84.8-86.0)	75.5 (74.8-76.1)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19	84.7 (75.0-90.9)	78.0 (68.3-85.0)	66.1 (57.5-73.4)
20-44	91.5 (89.0-93.5)	88.9 (86.2-91.1)	80.8 (77.8-83.4)
45-64	87.1 (84.4-89.3)	81.9 (79.1-84.4)	70.2 (67.4-72.8)
65+	80.8 (72.5-86.8)	74.4 (66.3-80.8)	64.1 (56.8-70.5)
Men	88.0 (85.7-89.9)	83.4 (81.0-85.5)	73.0 (70.5-75.3)
Women	89.1 (86.2-91.4)	85.4 (82.4-87.9)	75.8 (72.6-78.6)
Diabetes	88.3 (83.2-92.0)	82.8 (77.4-87.1)	69.9 (64.7-74.5)
Hypertension / renal vascular disease	78.3 (70.4-84.4)	71.1 (63.5-77.4)	61.4 (54.7-67.5)
Glomerulonephritis	90.5 (87.3-93.0)	87.4 (84.0-90.1)	79.4 (75.7-82.5)
Other cause	88.6 (86.1-90.7)	84.5 (81.8-86.8)	73.8 (71.1-76.4)
All	88.4 (86.7-89.9)	84.2 (82.3-85.8)	74.1 (72.1-75.9)

A5 SURVIVAL PROBABILITY
Graft Survival after First Transplant (cadaver donor)

Table A.5.26
One-, two- and five-year survival probabilities (cohort 1995-1999): First graft (cadaver donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	86.4 (83.5-89.3)	82.6 (79.4-85.9)	73.0 (69.3-76.9)
20-44	90.8 (90.0-91.7)	88.5 (87.6-89.4)	80.3 (79.1-81.4)
45-64	88.0 (87.1-88.8)	84.5 (83.5-85.5)	74.0 (72.8-75.2)
65+	81.3 (78.2-84.5)	75.4 (72.0-79.0)	57.8 (53.9-62.0)
Men	89.4 (88.7-90.2)	86.2 (85.4-87.0)	76.2 (75.2-77.2)
Women	88.0 (87.0-89.0)	85.0 (83.9-86.1)	75.5 (74.1-76.8)
Diabetes	86.5 (84.6-88.5)	83.0 (80.8-85.1)	70.0 (67.4-72.7)
Hypertension / renal vascular disease	87.7 (85.6-89.7)	83.0 (80.7-85.4)	72.3 (69.5-75.2)
Glomerulonephritis	90.2 (89.2-91.3)	87.5 (86.4-88.7)	77.6 (76.1-79.1)
Other cause	88.7 (87.8-89.5)	85.6 (84.7-86.5)	76.5 (75.4-77.6)
All	88.9 (88.3-89.5)	85.8 (85.1-86.4)	75.9 (75.1-76.8)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	91.8 (89.5-94.2)	89.3 (86.6-92.0)	81.3 (78.0-84.8)
45-64	87.1 (84.3-89.9)	82.1 (79.0-85.4)	70.6 (66.9-74.5)
65+	81.1 (73.0-90.1)	74.8 (65.9-84.9)	64.7 (55.1-76.0)
Men	88.5 (86.3-90.9)	84.1 (81.5-86.8)	74.0 (70.9-77.2)
Women	89.0 (86.2-91.9)	85.5 (82.4-88.8)	75.9 (72.1-79.9)
Diabetes	87.9 (83.0-93.0)	83.0 (77.5-88.9)	70.1 (63.4-77.4)
Hypertension / renal vascular disease	79.9 (72.1-88.6)	73.0 (64.4-82.8)	63.6 (54.3-74.4)
Glomerulonephritis	90.3 (87.2-93.5)	87.0 (83.5-90.7)	78.7 (74.5-83.2)
Other cause	89.0 (86.5-91.5)	85.0 (82.2-87.9)	74.6 (71.2-78.2)
All	88.7 (87.0-90.5)	84.6 (82.6-86.7)	74.7 (72.3-77.2)

Table A.5.27
One- and two-year survival probabilities (cohort 1998-2002): First graft (cadaver donor), unadjusted
from day of transplant to one and two years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	91.0 (88.5-93.1)	87.6 (84.9-89.9)
20-44	92.1 (91.4-92.8)	90.0 (89.2-90.7)
45-64	90.2 (89.5-90.9)	86.9 (86.2-87.7)
65+	84.6 (82.5-86.5)	78.5 (76.3-80.5)
Men	91.0 (90.4-91.5)	87.7 (87.1-88.3)
Women	90.0 (89.2-90.7)	87.3 (86.4-88.1)
Diabetes	90.5 (89.1-91.8)	86.9 (85.3-88.2)
Hypertension / renal vascular disease	87.4 (85.5-89.0)	83.1 (81.2-84.9)
Glomerulonephritis	91.8 (90.9-92.6)	88.9 (88.0-89.9)
Other cause	90.6 (89.9-91.2)	87.7 (87.0-88.4)
All	90.6 (90.1-91.1)	87.5 (87.0-88.0)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19	92.7 (80.5-97.4)	87.8 (75.5-94.2)
20-44	92.7 (90.4-94.5)	88.8 (86.2-90.9)
45-64	88.5 (86.1-90.5)	84.5 (82.0-86.8)
65+	82.0 (75.6-86.8)	77.0 (70.7-82.2)
Men	89.5 (87.5-91.2)	85.0 (82.8-86.9)
Women	90.2 (87.5-92.3)	86.8 (83.9-89.2)
Diabetes	90.2 (86.2-93.1)	86.5 (82.3-89.8)
Hypertension / renal vascular disease	82.6 (76.0-87.6)	77.4 (70.8-82.7)
Glomerulonephritis	91.3 (88.3-93.6)	87.0 (83.7-89.7)
Other cause	89.9 (87.5-91.9)	86.1 (83.5-88.3)
All	89.7 (88.2-91.1)	85.7 (84.0-87.2)

Table A.5.28
One- and two-year survival probabilities (cohort 1998-2002): First graft (cadaver donor), adjusted from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	90.7 (88.2-93.3)	87.2 (84.3-90.2)
20-44	92.2 (91.4-92.9)	90.1 (89.2-90.9)
45-64	90.3 (89.6-91.1)	87.1 (86.3-87.9)
65+	85.1 (82.8-87.4)	79.0 (76.4-81.7)
Men	91.4 (90.8-92.0)	88.4 (87.7-89.1)
Women	90.3 (89.5-91.1)	87.7 (86.7-88.6)
Diabetes	90.7 (89.2-92.1)	87.1 (85.4-88.7)
Hypertension / renal vascular disease	88.8 (87.0-90.6)	85.0 (83.0-87.1)
Glomerulonephritis	91.8 (90.9-92.7)	89.0 (88.0-90.1)
Other cause	90.9 (90.2-91.6)	88.2 (87.4-88.9)
All	91.0 (90.5-91.5)	88.1 (87.5-88.7)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	92.7 (90.5-94.9)	89.0 (86.4-91.7)
45-64	88.6 (86.2-91.1)	84.9 (82.2-87.7)
65+	82.5 (76.2-89.4)	77.7 (70.8-85.3)
Men	90.0 (88.1-92.0)	85.9 (83.7-88.2)
Women	90.5 (88.0-93.1)	87.5 (84.7-90.5)
Diabetes	90.2 (86.7-93.9)	87.1 (83.1-91.3)
Hypertension / renal vascular disease	84.4 (78.3-90.9)	79.6 (72.9-86.9)
Glomerulonephritis	91.1 (88.2-94.0)	87.0 (83.7-90.5)
Other cause	90.4 (88.2-92.7)	86.8 (84.2-89.4)
All	90.2 (88.7-91.8)	86.5 (84.7-88.3)

A5 SURVIVAL PROBABILITY
Graft Survival after First Transplant (living donor)

Table A.5.29
One-, two- and five-year survival probabilities (cohort 1995-1999): First graft (living donor), unadjusted
from day of transplant to one, two and five years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	93.4 (90.1-95.6)	90.6 (87.1-93.3)	81.9 (77.9-85.3)
20-44	94.6 (93.3-95.7)	92.9 (91.4-94.1)	86.1 (84.3-87.7)
45-64	92.6 (90.5-94.3)	91.7 (89.5-93.5)	85.4 (82.9-87.6)
65+	90.2 (80.8-95.1)	86.9 (77.4-92.6)	77.0 (67.7-84.0)
Men	94.6 (93.4-95.6)	93.2 (91.9-94.3)	85.6 (83.9-87.0)
Women	92.1 (90.2-93.7)	90.0 (87.9-91.7)	84.2 (82.0-86.3)
Diabetes	92.8 (88.3-95.6)	90.5 (85.8-93.7)	78.8 (73.6-83.1)
Hypertension / renal vascular disease	87.4 (81.5-91.5)	84.4 (78.4-88.9)	78.4 (72.4-83.3)
Glomerulonephritis	95.9 (94.3-97.1)	94.2 (92.3-95.6)	87.6 (85.3-89.5)
Other cause	93.2 (91.8-94.5)	91.9 (90.3-93.2)	85.3 (83.4-86.9)
All	93.7 (92.7-94.6)	92.1 (91.0-93.0)	85.1 (83.8-86.3)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44	96.9 (88.7-99.2)	95.4 (87.0-98.4)	86.2 (77.0-91.9)
45-64			
65+			
Men	94.1 (85.9-97.6)	92.6 (84.2-96.7)	86.7 (77.7-92.2)
Women	97.7 (85.2-99.7)	95.3 (83.6-98.7)	81.4 (69.6-89.0)
Diabetes			
Hypertension / renal vascular disease			
Glomerulonephritis	97.1 (82.3-99.6)	97.1 (82.3-99.6)	85.5 (71.8-92.9)
Other cause	95.3 (86.8-98.4)	92.2 (83.3-96.4)	84.3 (75.0-90.4)
All	95.5 (89.9-98.0)	93.7 (87.8-96.8)	84.6 (77.9-89.4)

A5 SURVIVAL PROBABILITY
Graft Survival after First Transplant (living donor)

Table A.5.30
One-, two- and five-year survival probabilities (cohort 1995-1999): First graft (living donor), adjusted from day of transplant to one, two and five years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
ALL COUNTRIES			
0-19	93.1 (90.1-96.1)	90.2 (86.7-93.8)	81.1 (76.5-85.9)
20-44	94.6 (93.3-95.9)	92.8 (91.4-94.3)	86.0 (84.1-88.0)
45-64	92.6 (90.6-94.7)	91.7 (89.6-93.8)	85.4 (82.7-88.2)
65+			
Men	94.4 (93.2-95.6)	92.9 (91.5-94.3)	84.9 (83.0-86.9)
Women	91.8 (89.8-93.8)	89.5 (87.3-91.8)	83.6 (80.9-86.3)
Diabetes	92.6 (88.8-96.5)	90.3 (86.0-94.8)	78.3 (72.5-84.7)
Hypertension / renal vascular disease	87.4 (82.0-93.2)	84.4 (78.6-90.8)	78.5 (71.8-85.7)
Glomerulonephritis	95.8 (94.3-97.3)	93.9 (92.2-95.7)	87.2 (84.7-89.7)
Other cause	92.9 (91.4-94.5)	91.5 (89.8-93.2)	84.6 (82.4-86.9)
All	93.5 (92.4-94.6)	91.7 (90.5-92.9)	84.4 (82.8-86.1)

	1 year Survival (95% CI)	2 year Survival (95% CI)	5 year Survival (95% CI)
AUSTRIA			
0-19			
20-44			
45-64			
65+			
Men			
Women			
Diabetes			
Hypertension / renal vascular disease			
Glomerulonephritis			
Other cause			
All			

Table A.5.31
One- and two-year survival probabilities (cohort 1998-2002): First graft (living donor), unadjusted
from day of transplant to one and two years, by age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	94.5 (91.6-96.4)	92.7 (89.5-94.9)
20-44	94.8 (93.6-95.8)	93.2 (91.9-94.3)
45-64	94.0 (92.3-95.3)	92.3 (90.5-93.7)
65+	87.0 (79.6-91.8)	84.8 (77.3-90.0)
Men	94.2 (93.1-95.2)	92.7 (91.5-93.7)
Women	94.3 (92.8-95.5)	92.3 (90.6-93.7)
Diabetes	92.6 (88.6-95.2)	90.2 (86.0-93.3)
Hypertension / renal vascular disease	93.2 (88.9-95.8)	91.1 (86.6-94.1)
Glomerulonephritis	95.4 (93.8-96.5)	94.3 (92.7-95.6)
Other cause	94.0 (92.8-95.0)	92.1 (90.7-93.2)
All	94.3 (93.4-95.0)	92.6 (91.6-93.4)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44	96.7 (90.6-98.9)	94.6 (88.0-97.6)
45-64	96.0 (85.7-98.9)	94.0 (83.6-97.9)
65+		
Men	95.1 (89.8-97.7)	92.7 (87.0-95.9)
Women	96.2 (86.2-99.0)	96.2 (86.2-99.0)
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis	93.5 (84.6-97.4)	91.9 (82.8-96.3)
Other cause	96.9 (91.1-99.0)	94.9 (88.7-97.7)
All	95.4 (91.3-97.6)	93.7 (89.3-96.3)

Table A.5.32
One- and two-year survival probabilities (cohort 1998-2002): First graft (living donor), adjusted
from day of transplant to one and two years, adjusted for age, gender and primary diagnosis

	1 year Survival (95% CI)	2 year Survival (95% CI)
ALL COUNTRIES		
0-19	94.4 (91.9-97.0)	92.5 (89.7-95.5)
20-44	94.8 (93.7-96.0)	93.2 (91.9-94.5)
45-64	94.1 (92.5-95.6)	92.4 (90.7-94.2)
65+		
Men	94.0 (92.9-95.1)	92.4 (91.1-93.7)
Women	94.2 (92.7-95.7)	92.2 (90.4-94.0)
Diabetes	92.5 (89.0-96.1)	90.1 (86.2-94.2)
Hypertension / renal vascular disease		
Glomerulonephritis	95.3 (93.8-96.7)	94.2 (92.6-95.8)
Other cause	93.7 (92.5-95.0)	91.8 (90.3-93.2)
All	94.1 (93.2-95.0)	92.3 (91.3-93.4)

	1 year Survival (95% CI)	2 year Survival (95% CI)
AUSTRIA		
0-19		
20-44		
45-64		
65+		
Men		
Women		
Diabetes		
Hypertension / renal vascular disease		
Glomerulonephritis		
Other cause		
All		

ERA-EDTA Registry
Academic Medical Center
Department of Medical Informatics
Room J1B-125
PO Box 22700
1100 DE Amsterdam
The Netherlands

www.era-edta-reg.org
July 2006