



Leading European Nephrology

Austria

Survival
Report
2015

ERA-EDTA Registry

Academic Medical Center
Department of Medical Informatics
PO Box 22700
1100 DE Amsterdam
the Netherlands

Telephone +31 20 566 7637
Fax +31 20 566 9014
E-mail erareg@amc.uva.nl
Website www.era-edta-reg.org

Further information (e.g. collaborating registries, methods) can be found in the ERA-EDTA Registry annual report 2015, which is published on the ERA-EDTA Registry website www.era-edta-reg.org.

ERA-EDTA Registry
Survival Report 2015
Austria

Table of Contents

A5 SURVIVAL PROBABILITY

Table A.5.1 One-, two- and five-year survival probabilities by treatment modality and cohort	6
--	---

B6 SURVIVAL PROBABILITY

Patient Survival on RRT (cohort 2006-2010 and cohort 2009-2013)

Table B.6.1 Incident RRT patients, from day 1, unadjusted	8
Table B.6.2 Incident RRT patients, from day 1, adjusted	10
Table B.6.3 Incident RRT patients, from day 91, unadjusted	12
Table B.6.4 Incident RRT patients, from day 91, adjusted	14

Patient Survival on Dialysis (cohort 2006-2010 and cohort 2009-2013)

Table B.6.5 Incident dialysis patients, from day 1, unadjusted	16
Table B.6.6 Incident dialysis patients, from day 1, adjusted	18
Table B.6.7 Incident dialysis patients, from day 91, unadjusted	20
Table B.6.8 Incident dialysis patients, from day 91, adjusted	22

Patient Survival after First Transplant (cohort 2006-2010 and cohort 2009-2013)

Table B.6.9 First transplant patients (deceased donor), from day of transplant, unadjusted	24
Table B.6.10 First transplant patients (deceased donor), from day of transplant, adjusted	26
Table B.6.11 First transplant patients (living donor), from day of transplant, unadjusted	28
Table B.6.12 First transplant patients (living donor), from day of transplant, adjusted	30

Graft Survival after First Transplant (cohort 2006-2010 and cohort 2009-2013)

Table B.6.13 First graft (deceased donor), from day of transplant, unadjusted	32
Table B.6.14 First graft (deceased donor), from day of transplant, adjusted	34
Table B.6.15 First graft (living donor), from day of transplant, unadjusted	36
Table B.6.16 First graft (living donor), from day of transplant, adjusted	38

Table A.5.1

One-, two- and five-year survival probabilities by treatment modality and cohort*from day 1 of RRT / dialysis, or from day of transplant*

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
Patient survival on RRT					
Unadjusted	83.2 (83.1-83.4)	72.7 (72.6-72.9)	50.0 (49.9-50.1)	84.4 (84.2-84.5)	74.3 (74.2-74.5)
Adjusted †	85.9 (85.8-86.1)	76.1 (75.9-76.3)	51.1 (50.8-51.4)	86.9 (86.7-87.1)	77.6 (77.4-77.9)
Patient survival on dialysis					
Unadjusted	82.2 (82.1-82.4)	70.4 (70.2-70.6)	41.8 (41.7-41.9)	83.3 (83.1-83.4)	71.9 (71.7-72.0)
Adjusted †	84.4 (84.2-84.6)	73.5 (73.3-73.7)	45.0 (44.7-45.4)	85.7 (85.5-85.9)	75.4 (75.2-75.6)
Patient survival after first kidney transplantation (deceased donor)					
Unadjusted	96.1 (95.9-96.4)	94.3 (94.1-94.6)	88.0 (87.6-88.3)	96.3 (96.1-96.5)	94.3 (94.0-94.5)
Adjusted ††	97.5 (97.3-97.6)	96.2 (96.0-96.4)	91.7 (91.3-92.1)	97.8 (97.7-98.0)	96.5 (96.3-96.7)
Graft survival after first kidney transplantation (deceased donor)					
Unadjusted	90.9 (90.5-91.2)	88.1 (87.8-88.5)	78.9 (78.5-79.3)	91.1 (90.8-91.3)	88.1 (87.7-88.4)
Adjusted ††	91.8 (91.4-92.1)	89.3 (88.9-89.7)	80.7 (80.2-81.2)	92.4 (92.1-92.7)	89.8 (89.4-90.1)
Patient survival after first kidney transplantation (living donor)					
Unadjusted	98.6 (98.3-98.8)	97.7 (97.4-98.0)	94.1 (93.6-94.6)	98.9 (98.7-99.1)	97.9 (97.6-98.2)
Adjusted ††	98.7 (98.5-99.0)	98.0 (97.7-98.3)	94.5 (94.0-95.0)	99.1 (98.9-99.2)	98.2 (98.0-98.5)
Graft survival after first kidney transplantation (living donor)					
Unadjusted	95.9 (95.4-96.3)	94.1 (93.6-94.5)	87.4 (86.7-88.0)	96.5 (96.2-96.8)	94.7 (94.3-95.1)
Adjusted ††	95.6 (95.2-96.1)	93.8 (93.2-94.3)	86.6 (85.8-87.4)	96.3 (95.9-96.7)	94.4 (93.9-94.9)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

† Analyses were adjusted using fixed values: age (67 years), gender (63% men) and PRD (24% diabetes mellitus, 19% hypertension/renal vascular disease, 11% glomerulonephritis and 46% other causes).

†† Analyses were adjusted using fixed values: age (50 years), gender (63% men) and PRD (14% diabetes mellitus, 10% hypertension/renal vascular disease, 23% glomerulonephritis and 53% other causes).

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
Patient survival on RRT					
Unadjusted	79.7 (78.8-80.5)	67.2 (66.4-68.0)	43.7 (43.1-44.2)	82.2 (81.4-83.0)	70.7 (69.9-71.5)
Adjusted †	82.0 (81.1-82.9)	69.8 (68.7-71.0)	42.6 (41.3-44.0)	84.3 (83.4-85.2)	73.2 (72.1-74.3)
Patient survival on dialysis					
Unadjusted	78.9 (78.1-79.7)	65.1 (64.3-65.9)	36.3 (35.8-36.8)	81.5 (80.7-82.3)	68.8 (67.9-69.6)
Adjusted †	80.5 (79.5-81.5)	67.1 (65.9-68.3)	36.8 (35.4-38.2)	83.1 (82.2-84.1)	71.0 (69.9-72.2)
Patient survival after first kidney transplantation (deceased donor)					
Unadjusted	93.2 (91.7-94.4)	90.3 (88.7-91.7)	83.6 (81.8-85.2)	95.0 (93.7-96.0)	92.7 (91.2-93.9)
Adjusted ††	96.0 (95.1-96.8)	94.1 (93.1-95.2)	89.8 (88.5-91.2)	97.3 (96.6-97.9)	96.0 (95.2-96.8)
Graft survival after first kidney transplantation (deceased donor)					
Unadjusted	88.4 (86.8-89.9)	84.4 (82.6-86.0)	75.3 (73.5-77.1)	90.1 (88.5-91.4)	85.9 (84.2-87.5)
Adjusted ††	90.2 (88.7-91.7)	86.6 (84.9-88.4)	78.6 (76.5-80.8)	92.0 (90.6-93.3)	88.5 (87.0-90.1)
Patient survival after first kidney transplantation (living donor)					
Unadjusted	98.8 (96.5-99.6)	98.1 (95.5-99.2)	94.5 (91.2-96.6)	98.9 (96.7-99.6)	98.5 (96.2-99.4)
Adjusted ††	98.8 (97.5-100.0)	98.0 (96.3-99.8)	94.2 (91.4-97.2)	98.9 (97.7-100.0)	98.5 (97.1-100.0)
Graft survival after first kidney transplantation (living donor)					
Unadjusted	97.3 (94.5-98.7)	95.3 (92.2-97.3)	88.3 (84.3-91.3)	96.3 (93.4-98.0)	95.6 (92.6-97.4)
Adjusted ††	97.0 (94.8-99.2)	94.8 (92.0-97.7)	87.0 (82.7-91.4)	95.9 (93.5-98.4)	95.1 (92.4-97.8)

† Analyses were adjusted using fixed values: age (67 years), gender (63% men) and PRD (24% diabetes mellitus, 19% hypertension/renal vascular disease, 11% glomerulonephritis and 46% other causes).

†† Analyses were adjusted using fixed values: age (50 years), gender (63% men) and PRD (14% diabetes mellitus, 10% hypertension/renal vascular disease, 23% glomerulonephritis and 53% other causes).

Table B.6.1
Incident RRT patients, from day 1, unadjusted
by age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)						
	cohort 2006 - 2010				cohort 2009 - 2013		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.2 (98.7-99.6)	97.1 (96.2-97.8)	96.1 (95.1-97.0)	94.4 (93.2-95.4)	99.3 (98.8-99.6)	97.7 (96.9-98.4)	96.4 (95.4-97.2)
20-44	99.1 (99.0-99.2)	96.9 (96.7-97.2)	94.4 (94.1-94.7)	89.0 (88.5-89.4)	99.2 (99.1-99.3)	97.2 (97.0-97.5)	94.9 (94.6-95.2)
45-64	96.9 (96.8-97.1)	90.5 (90.3-90.7)	83.7 (83.4-84.0)	67.2 (66.9-67.5)	97.3 (97.2-97.4)	91.3 (91.1-91.5)	85.2 (85.0-85.5)
65-74	93.7 (93.5-94.0)	81.5 (81.2-81.8)	69.6 (69.3-69.9)	42.3 (42.1-42.5)	94.4 (94.2-94.6)	83.5 (83.2-83.8)	72.4 (72.1-72.7)
75+	89.7 (89.4-89.9)	72.0 (71.7-72.3)	55.7 (55.5-56.0)	23.4 (23.3-23.5)	90.5 (90.3-90.7)	73.8 (73.6-74.1)	58.1 (57.9-58.4)
Men	94.2 (94.1-94.4)	83.2 (83.0-83.4)	72.5 (72.3-72.7)	49.3 (49.2-49.5)	94.6 (94.4-94.7)	84.3 (84.1-84.5)	74.1 (73.9-74.3)
Women	93.9 (93.8-94.1)	83.3 (83.0-83.5)	73.1 (72.8-73.3)	51.2 (51.0-51.4)	94.6 (94.4-94.8)	84.5 (84.2-84.7)	74.9 (74.6-75.1)
Diabetes	94.9 (94.7-95.1)	83.6 (83.3-84.0)	71.0 (70.6-71.3)	42.6 (42.4-42.9)	95.4 (95.2-95.6)	85.1 (84.8-85.4)	73.5 (73.2-73.8)
Hypertension / renal vascular disease	94.1 (93.8-94.3)	81.6 (81.3-82.0)	68.6 (68.2-69.0)	40.4 (40.2-40.7)	94.2 (94.0-94.5)	82.8 (82.4-83.1)	70.2 (69.8-70.5)
Glomerulonephritis	97.4 (97.1-97.6)	92.0 (91.6-92.4)	86.3 (85.9-86.8)	71.9 (71.4-72.3)	97.3 (97.1-97.6)	92.2 (91.8-92.5)	86.8 (86.4-87.2)
Other causes	93.3 (93.1-93.5)	82.2 (81.9-82.4)	72.7 (72.4-72.9)	52.8 (52.6-53.0)	93.9 (93.8-94.1)	83.2 (82.9-83.4)	73.9 (73.6-74.1)
All	94.1 (94.0-94.2)	83.2 (83.1-83.4)	72.7 (72.6-72.9)	50.0 (49.9-50.1)	94.6 (94.5-94.7)	84.4 (84.2-84.5)	74.3 (74.2-74.5)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)						
	cohort 2006 - 2010				cohort 2009 - 2013		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19							
20-44	98.5 (97.1-99.2)	95.9 (94.1-97.2)	93.1 (90.9-94.8)	88.6 (86.0-90.7)	99.3 (98.1-99.7)	96.1 (94.2-97.4)	93.7 (91.5-95.4)
45-64	95.1 (94.1-95.9)	87.7 (86.3-88.9)	79.8 (78.3-81.2)	60.2 (58.9-61.5)	96.3 (95.4-97.0)	89.5 (88.2-90.7)	82.6 (81.1-84.0)
65-74	92.0 (90.8-93.1)	76.7 (75.1-78.2)	62.0 (60.6-63.4)	33.7 (32.9-34.4)	93.6 (92.4-94.5)	79.4 (77.9-80.9)	65.5 (64.0-67.0)
75+	87.7 (86.3-89.0)	66.6 (65.0-68.1)	46.9 (45.7-48.0)	15.5 (15.2-15.7)	90.7 (89.3-91.9)	71.4 (69.8-73.0)	53.6 (52.2-54.9)
Men	93.1 (92.3-93.8)	80.3 (79.3-81.3)	67.7 (66.7-68.7)	43.6 (42.9-44.3)	94.5 (93.8-95.1)	82.8 (81.8-83.8)	71.3 (70.3-72.3)
Women	91.7 (90.6-92.7)	78.6 (77.2-79.9)	66.4 (65.1-67.7)	43.7 (42.8-44.6)	93.8 (92.7-94.7)	81.2 (79.8-82.5)	69.6 (68.2-71.0)
Diabetes	93.3 (92.2-94.3)	78.9 (77.4-80.3)	63.0 (61.6-64.3)	35.3 (34.5-36.0)	95.4 (94.3-96.2)	82.0 (80.5-83.5)	67.6 (66.1-69.1)
Hypertension / renal vascular disease	93.2 (91.9-94.4)	78.2 (76.4-79.9)	63.0 (61.3-64.6)	36.1 (35.1-37.0)	94.4 (93.2-95.5)	81.1 (79.3-82.7)	67.8 (66.1-69.5)
Glomerulonephritis	97.0 (95.5-98.0)	91.9 (89.8-93.6)	85.6 (83.1-87.7)	67.4 (64.9-69.8)	97.0 (95.4-98.1)	93.2 (91.1-94.8)	88.0 (85.5-90.0)
Other causes	90.1 (88.9-91.2)	77.3 (75.9-78.7)	67.8 (66.4-69.2)	48.5 (47.4-49.6)	92.5 (91.4-93.4)	79.9 (78.6-81.3)	69.9 (68.5-71.3)
All	92.6 (91.9-93.2)	79.7 (78.8-80.5)	67.2 (66.4-68.0)	43.7 (43.1-44.2)	94.2 (93.7-94.8)	82.2 (81.4-83.0)	70.7 (69.9-71.5)

Table B.6.2
Incident RRT patients, from day 1, adjusted
adjusted for age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)						
	cohort 2006 - 2010				cohort 2009 - 2013		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.1 (98.6-99.6)	96.9 (95.9-97.8)	95.9 (94.8-96.9)	94.0 (92.8-95.2)	99.3 (98.9-99.7)	97.8 (97.0-98.5)	96.4 (95.4-97.4)
20-44	99.1 (99.0-99.2)	96.9 (96.6-97.1)	94.3 (93.9-94.6)	88.5 (88.1-89.0)	99.2 (99.1-99.3)	97.2 (97.0-97.5)	94.7 (94.4-95.1)
45-64	97.1 (96.9-97.2)	90.8 (90.6-91.1)	84.1 (83.8-84.5)	67.5 (67.1-68.0)	97.4 (97.3-97.6)	91.5 (91.3-91.8)	85.5 (85.2-85.8)
65-74	94.1 (93.9-94.4)	82.5 (82.1-82.9)	70.8 (70.4-71.3)	43.5 (43.0-44.0)	94.7 (94.5-94.9)	84.2 (83.8-84.5)	73.2 (72.8-73.7)
75+	89.9 (89.7-90.2)	72.7 (72.3-73.1)	56.5 (56.1-57.0)	23.8 (23.4-24.2)	90.7 (90.5-91.0)	74.4 (74.0-74.8)	58.8 (58.4-59.3)
Men	95.3 (95.2-95.4)	85.7 (85.5-86.0)	75.6 (75.3-75.9)	49.8 (49.4-50.1)	95.6 (95.5-95.7)	86.7 (86.5-86.9)	77.2 (76.9-77.4)
Women	95.2 (95.1-95.4)	86.3 (86.0-86.5)	76.9 (76.6-77.3)	53.3 (52.8-53.7)	95.7 (95.5-95.8)	87.2 (86.9-87.5)	78.4 (78.1-78.7)
Diabetes	95.4 (95.2-95.6)	85.0 (84.6-85.3)	72.6 (72.2-73.1)	42.1 (41.6-42.7)	95.9 (95.7-96.1)	86.6 (86.3-86.9)	75.5 (75.1-76.0)
Hypertension / renal vascular disease	96.2 (96.1-96.4)	88.0 (87.7-88.3)	78.4 (77.9-78.8)	52.6 (52.0-53.2)	96.3 (96.2-96.5)	88.7 (88.4-89.0)	79.5 (79.1-79.9)
Glomerulonephritis	96.8 (96.6-97.1)	90.3 (89.9-90.8)	83.1 (82.5-83.8)	63.4 (62.6-64.3)	96.9 (96.6-97.1)	90.6 (90.2-91.1)	83.9 (83.3-84.5)
Other causes	94.1 (94.0-94.3)	83.9 (83.6-84.2)	74.5 (74.1-74.8)	51.7 (51.3-52.1)	94.7 (94.6-94.9)	85.0 (84.7-85.2)	75.9 (75.6-76.2)
All	95.3 (95.2-95.4)	85.9 (85.8-86.1)	76.1 (75.9-76.3)	51.1 (50.8-51.4)	95.6 (95.5-95.7)	86.9 (86.7-87.1)	77.6 (77.4-77.9)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)						
	cohort 2006 - 2010				cohort 2009 - 2013		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19							
20-44	98.4 (97.4-99.4)	95.3 (93.6-97.1)	92.8 (90.6-94.9)	87.9 (85.2-90.7)	99.2 (98.5-100.0)	95.5 (93.8-97.3)	93.4 (91.2-95.5)
45-64	95.2 (94.3-96.1)	87.9 (86.5-89.4)	80.2 (78.5-82.0)	60.8 (58.7-63.0)	96.2 (95.4-97.1)	89.6 (88.2-91.0)	82.8 (81.0-84.5)
65-74	92.4 (91.3-93.6)	77.7 (75.9-79.7)	63.6 (61.4-65.9)	35.4 (33.2-37.8)	93.8 (92.7-94.9)	79.9 (78.1-81.8)	66.3 (64.1-68.5)
75+	87.9 (86.4-89.5)	67.4 (65.2-69.7)	48.1 (45.7-50.6)	16.5 (14.7-18.4)	90.7 (89.3-92.1)	71.9 (69.7-74.1)	54.1 (51.8-56.7)
Men	93.8 (93.1-94.6)	81.8 (80.7-83.0)	69.1 (67.6-70.6)	40.9 (39.2-42.6)	95.1 (94.4-95.7)	84.3 (83.1-85.4)	72.9 (71.5-74.3)
Women	93.5 (92.6-94.4)	82.4 (81.0-83.9)	71.1 (69.4-72.9)	45.6 (43.5-47.9)	95.0 (94.1-95.8)	84.3 (82.9-85.8)	73.8 (72.0-75.6)
Diabetes	94.1 (93.1-95.1)	80.6 (79.0-82.4)	65.0 (62.9-67.1)	34.5 (32.4-36.9)	95.8 (95.0-96.7)	83.6 (82.0-85.4)	69.9 (67.8-72.1)
Hypertension / renal vascular disease	95.1 (94.2-96.1)	84.2 (82.5-85.9)	71.6 (69.4-73.8)	44.2 (41.6-47.0)	96.1 (95.2-96.9)	86.4 (84.8-87.9)	75.6 (73.7-77.7)
Glomerulonephritis	96.0 (94.3-97.7)	88.9 (86.2-91.7)	79.8 (76.3-83.5)	53.2 (48.8-58.0)	96.1 (94.3-97.8)	90.9 (88.3-93.5)	83.6 (80.2-87.0)
Other causes	90.7 (89.5-91.9)	77.7 (75.9-79.5)	67.5 (65.4-69.5)	43.0 (40.7-45.5)	93.0 (91.9-94.0)	80.6 (79.0-82.3)	70.2 (68.3-72.2)
All	93.7 (93.1-94.2)	82.0 (81.1-82.9)	69.8 (68.7-71.0)	42.6 (41.3-44.0)	95.0 (94.5-95.5)	84.3 (83.4-85.2)	73.2 (72.1-74.3)

Table B.6.3
Incident RRT patients, from day 91, unadjusted
by age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	97.6 (96.7-98.2)	96.7 (95.7-97.4)	95.1 (93.9-96.0)	98.0 (97.1-98.6)	97.0 (96.0-97.7)
20-44	97.2 (97.0-97.5)	94.7 (94.4-95.0)	89.3 (88.8-89.7)	97.4 (97.2-97.6)	95.2 (94.9-95.5)
45-64	91.5 (91.3-91.8)	84.9 (84.6-85.1)	67.9 (67.6-68.2)	92.3 (92.1-92.5)	86.1 (85.8-86.3)
65-74	83.7 (83.3-84.0)	71.5 (71.1-71.8)	42.9 (42.7-43.1)	85.6 (85.3-85.9)	73.9 (73.5-74.2)
75+	75.7 (75.4-76.0)	58.2 (57.9-58.5)	23.9 (23.8-24.0)	77.3 (77.0-77.6)	60.3 (60.0-60.5)
Men	85.4 (85.2-85.6)	74.5 (74.3-74.8)	50.6 (50.4-50.7)	86.4 (86.3-86.6)	75.8 (75.6-76.0)
Women	85.9 (85.6-86.2)	75.5 (75.2-75.7)	52.8 (52.5-53.0)	86.9 (86.6-87.1)	76.8 (76.5-77.1)
Diabetes	84.7 (84.3-85.0)	71.8 (71.5-72.2)	42.7 (42.4-42.9)	86.1 (85.8-86.4)	74.0 (73.7-74.4)
Hypertension / renal vascular disease	83.3 (82.9-83.6)	69.8 (69.4-70.2)	40.9 (40.6-41.1)	84.6 (84.2-84.9)	71.5 (71.1-71.9)
Glomerulonephritis	93.1 (92.7-93.5)	87.2 (86.8-87.7)	72.6 (72.1-73.1)	93.4 (93.0-93.7)	87.6 (87.2-88.0)
Other causes	85.5 (85.2-85.7)	75.8 (75.5-76.0)	55.1 (54.8-55.3)	86.1 (85.9-86.4)	76.4 (76.2-76.7)
All	85.6 (85.4-85.7)	74.9 (74.7-75.1)	51.4 (51.2-51.5)	86.6 (86.4-86.7)	76.1 (76.0-76.3)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	96.0 (94.1-97.3)	94.4 (92.3-95.9)	89.6 (87.1-91.7)	95.5 (93.5-96.9)	94.1 (91.9-95.8)
45-64	90.1 (88.8-91.2)	82.4 (80.9-83.8)	61.5 (60.1-62.9)	91.2 (89.9-92.4)	84.3 (82.8-85.7)
65-74	79.4 (77.8-80.9)	63.6 (62.1-65.1)	34.6 (33.8-35.5)	80.9 (79.3-82.4)	66.5 (65.0-68.1)
75+	69.8 (68.1-71.4)	49.4 (48.1-50.7)	15.8 (15.5-16.1)	73.5 (71.8-75.2)	55.0 (53.5-56.4)
Men	82.7 (81.6-83.7)	70.1 (69.1-71.2)	45.2 (44.4-46.0)	84.4 (83.3-85.4)	72.9 (71.8-74.0)
Women	82.3 (80.9-83.6)	69.8 (68.3-71.1)	46.0 (45.0-47.0)	83.3 (81.8-84.6)	71.5 (70.0-73.0)
Diabetes	80.1 (78.5-81.5)	64.6 (63.1-66.0)	35.3 (34.5-36.1)	81.5 (79.8-83.0)	68.0 (66.4-69.5)
Hypertension / renal vascular disease	80.0 (78.1-81.8)	64.5 (62.7-66.2)	37.1 (36.1-38.2)	82.6 (80.8-84.2)	69.1 (67.3-70.8)
Glomerulonephritis	92.6 (90.5-94.3)	86.4 (83.9-88.5)	68.8 (66.2-71.2)	94.0 (92.0-95.6)	88.2 (85.7-90.3)
Other causes	82.9 (81.4-84.3)	73.0 (71.5-74.5)	52.6 (51.4-53.8)	83.8 (82.4-85.2)	73.3 (71.8-74.7)
All	82.5 (81.7-83.3)	70.0 (69.1-70.9)	45.5 (44.9-46.1)	84.0 (83.2-84.8)	72.4 (71.6-73.3)

Table B.6.4
Incident RRT patients, from day 91, adjusted
adjusted for age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	97.4 (96.5-98.2)	96.4 (95.4-97.3)	94.6 (93.5-95.8)	98.0 (97.2-98.7)	96.9 (96.0-97.8)
20-44	97.1 (96.9-97.4)	94.5 (94.2-94.9)	88.9 (88.4-89.4)	97.3 (97.1-97.6)	95.0 (94.7-95.4)
45-64	91.8 (91.5-92.0)	85.2 (84.9-85.5)	68.4 (68.0-68.9)	92.4 (92.2-92.6)	86.3 (86.0-86.6)
65-74	84.4 (84.0-84.8)	72.6 (72.1-73.1)	44.6 (44.1-45.1)	86.0 (85.7-86.4)	74.8 (74.4-75.3)
75+	76.2 (75.8-76.6)	58.9 (58.4-59.4)	24.7 (24.3-25.2)	77.7 (77.4-78.1)	61.1 (60.7-61.6)
Men	87.3 (87.1-87.5)	76.9 (76.6-77.2)	50.6 (50.3-51.0)	88.3 (88.1-88.5)	78.4 (78.1-78.7)
Women	88.2 (87.9-88.5)	78.6 (78.2-78.9)	54.4 (53.9-54.8)	88.9 (88.6-89.2)	79.8 (79.4-80.1)
Diabetes	85.7 (85.3-86.1)	73.1 (72.7-73.6)	42.1 (41.5-42.6)	87.3 (87.0-87.7)	75.9 (75.5-76.4)
Hypertension / renal vascular disease	88.9 (88.6-89.3)	79.0 (78.6-79.5)	52.9 (52.3-53.5)	89.8 (89.5-90.1)	80.3 (79.9-80.8)
Glomerulonephritis	91.5 (91.0-92.0)	83.9 (83.3-84.6)	64.1 (63.3-65.0)	91.9 (91.4-92.3)	84.7 (84.1-85.3)
Other causes	86.6 (86.3-86.9)	76.9 (76.5-77.2)	53.4 (53.0-53.8)	87.3 (87.1-87.6)	77.9 (77.6-78.3)
All	87.6 (87.5-87.8)	77.5 (77.3-77.8)	52.0 (51.7-52.3)	88.5 (88.4-88.7)	78.9 (78.7-79.2)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	95.8 (94.1-97.5)	94.1 (92.1-96.1)	89.1 (86.5-91.8)	95.2 (93.4-97.1)	93.8 (91.7-95.9)
45-64	90.4 (89.1-91.8)	83.0 (81.3-84.7)	62.7 (60.5-65.0)	91.4 (90.1-92.7)	84.7 (83.0-86.4)
65-74	80.5 (78.6-82.4)	65.5 (63.2-67.8)	37.2 (34.9-39.7)	81.6 (79.7-83.5)	67.8 (65.5-70.1)
75+	70.9 (68.6-73.3)	50.9 (48.4-53.6)	17.4 (15.4-19.5)	74.1 (71.9-76.4)	56.1 (53.6-58.8)
Men	83.9 (82.8-85.1)	71.1 (69.6-72.6)	42.4 (40.6-44.2)	85.6 (84.5-86.7)	74.5 (73.0-75.9)
Women	85.3 (83.9-86.7)	73.7 (71.9-75.6)	47.3 (45.1-49.7)	86.0 (84.6-87.5)	75.3 (73.4-77.1)
Diabetes	81.6 (79.8-83.3)	66.2 (64.0-68.4)	34.4 (32.1-36.8)	83.1 (81.3-84.9)	70.2 (68.0-72.4)
Hypertension / renal vascular disease	85.3 (83.6-87.1)	72.3 (70.1-74.6)	45.1 (42.3-48.0)	87.3 (85.8-88.9)	76.6 (74.6-78.7)
Glomerulonephritis	89.5 (86.8-92.4)	80.4 (76.8-84.2)	54.9 (50.4-59.8)	91.9 (89.4-94.5)	83.9 (80.5-87.4)
Other causes	82.7 (81.0-84.5)	71.8 (69.7-73.9)	46.0 (43.6-48.7)	84.2 (82.6-85.8)	73.2 (71.2-75.2)
All	84.4 (83.5-85.4)	72.1 (70.9-73.3)	44.2 (42.8-45.6)	85.8 (84.9-86.7)	74.8 (73.6-75.9)

Table B.6.5
Incident dialysis patients, from day 1, unadjusted
by age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)						
	cohort 2006 - 2010				cohort 2009 - 2013		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	99.0 (98.2-99.4)	96.1 (94.7-97.1)	94.1 (92.3-95.6)	88.8 (85.5-91.4)	99.2 (98.5-99.6)	97.1 (95.8-98.0)	94.0 (92.0-95.5)
20-44	99.0 (98.9-99.2)	96.4 (96.0-96.6)	92.7 (92.3-93.1)	82.2 (81.5-82.9)	99.1 (99.0-99.3)	96.7 (96.4-97.0)	93.2 (92.8-93.6)
45-64	96.8 (96.6-97.0)	89.8 (89.5-90.0)	81.7 (81.4-82.1)	59.0 (58.7-59.3)	97.2 (97.0-97.3)	90.5 (90.2-90.7)	83.1 (82.8-83.4)
65-74	93.6 (93.4-93.9)	81.1 (80.8-81.4)	68.8 (68.4-69.1)	39.5 (39.3-39.7)	94.2 (94.0-94.5)	83.0 (82.6-83.3)	71.3 (70.9-71.6)
75+	89.6 (89.3-89.8)	71.9 (71.6-72.2)	55.6 (55.3-55.8)	23.1 (23.0-23.2)	90.4 (90.2-90.7)	73.7 (73.5-74.0)	57.9 (57.7-58.2)
Men	94.0 (93.8-94.1)	82.2 (82.0-82.4)	70.2 (70.0-70.4)	41.0 (40.9-41.2)	94.3 (94.2-94.5)	83.3 (83.1-83.5)	71.6 (71.4-71.8)
Women	93.7 (93.5-93.9)	82.2 (81.9-82.5)	70.7 (70.4-71.0)	43.1 (42.9-43.3)	94.3 (94.1-94.5)	83.3 (83.1-83.6)	72.3 (72.0-72.6)
Diabetes	94.8 (94.5-95.0)	83.1 (82.7-83.4)	69.6 (69.3-70.0)	37.4 (37.2-37.6)	95.3 (95.1-95.5)	84.7 (84.3-85.0)	72.3 (71.9-72.6)
Hypertension / renal vascular disease	93.9 (93.7-94.2)	81.2 (80.8-81.6)	67.5 (67.1-67.9)	36.1 (35.8-36.3)	94.1 (93.9-94.4)	82.3 (81.9-82.7)	69.0 (68.6-69.3)
Glomerulonephritis	97.2 (96.9-97.4)	91.1 (90.7-91.5)	83.9 (83.3-84.4)	61.4 (60.8-61.9)	97.1 (96.9-97.4)	91.2 (90.7-91.6)	84.2 (83.7-84.7)
Other causes	93.0 (92.8-93.1)	80.9 (80.6-81.1)	69.9 (69.6-70.1)	43.7 (43.5-43.9)	93.5 (93.4-93.7)	81.7 (81.5-81.9)	70.8 (70.5-71.0)
All	93.9 (93.7-94.0)	82.2 (82.1-82.4)	70.4 (70.2-70.6)	41.8 (41.7-41.9)	94.3 (94.2-94.4)	83.3 (83.1-83.4)	71.9 (71.7-72.0)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)						
	cohort 2006 - 2010				cohort 2009 - 2013		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19							
20-44	98.7 (97.2-99.4)	95.8 (93.7-97.3)	91.7 (88.9-93.8)	83.5 (79.8-86.7)	99.2 (97.9-99.7)	95.7 (93.5-97.2)	92.1 (89.3-94.2)
45-64	95.0 (94.0-95.8)	87.3 (85.9-88.5)	78.4 (76.8-79.9)	53.0 (51.6-54.4)	96.2 (95.2-96.9)	89.1 (87.7-90.3)	81.2 (79.6-82.7)
65-74	92.0 (90.8-93.1)	76.5 (74.9-78.0)	61.1 (59.7-62.5)	31.2 (30.4-31.9)	93.6 (92.4-94.6)	79.2 (77.7-80.7)	64.7 (63.2-66.1)
75+	87.7 (86.3-89.0)	66.6 (65.0-68.1)	46.9 (45.7-48.0)	15.3 (15.1-15.6)	90.7 (89.3-91.9)	71.4 (69.8-72.9)	53.5 (52.1-54.8)
Men	92.9 (92.1-93.6)	79.6 (78.5-80.6)	65.5 (64.4-66.5)	35.9 (35.3-36.6)	94.4 (93.6-95.0)	82.1 (81.1-83.1)	69.4 (68.3-70.4)
Women	91.6 (90.5-92.6)	77.8 (76.4-79.1)	64.5 (63.1-65.8)	36.9 (36.1-37.7)	93.6 (92.6-94.6)	80.4 (79.0-81.8)	67.7 (66.2-69.1)
Diabetes	93.3 (92.1-94.2)	78.5 (77.0-79.9)	61.4 (60.0-62.8)	30.3 (29.6-31.0)	95.3 (94.3-96.2)	81.8 (80.2-83.3)	66.3 (64.7-67.8)
Hypertension / renal vascular disease	93.2 (91.8-94.4)	78.1 (76.3-79.8)	62.2 (60.5-63.8)	32.1 (31.2-33.0)	94.4 (93.1-95.4)	80.9 (79.1-82.5)	67.1 (65.3-68.7)
Glomerulonephritis	96.8 (95.3-97.9)	91.3 (89.0-93.1)	83.4 (80.7-85.8)	58.3 (55.6-61.0)	96.8 (95.1-97.9)	92.6 (90.3-94.4)	86.0 (83.2-88.3)
Other causes	89.7 (88.5-90.8)	76.0 (74.5-77.4)	65.1 (63.7-66.5)	39.4 (38.4-40.3)	92.2 (91.1-93.2)	78.8 (77.3-80.1)	67.4 (66.0-68.8)
All	92.4 (91.8-93.0)	78.9 (78.1-79.7)	65.1 (64.3-65.9)	36.3 (35.8-36.8)	94.1 (93.5-94.7)	81.5 (80.7-82.3)	68.8 (67.9-69.6)

Table B.6.6
Incident dialysis patients, from day 1, adjusted
adjusted for age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)						
	cohort 2006 - 2010				cohort 2009 - 2013		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19	98.8 (98.2-99.5)	95.8 (94.4-97.2)	93.8 (91.9-95.7)	86.7 (82.5-91.2)	99.1 (98.5-99.7)	97.0 (95.8-98.1)	93.9 (91.9-95.8)
20-44	99.0 (98.9-99.2)	96.4 (96.0-96.7)	92.6 (92.1-93.1)	80.9 (80.0-81.8)	99.1 (99.0-99.3)	96.7 (96.4-97.0)	93.1 (92.7-93.6)
45-64	96.9 (96.8-97.1)	90.1 (89.9-90.4)	82.3 (81.9-82.7)	58.7 (58.1-59.2)	97.3 (97.1-97.4)	90.8 (90.5-91.0)	83.5 (83.1-83.9)
65-74	94.0 (93.7-94.2)	82.0 (81.6-82.4)	69.9 (69.4-70.4)	40.1 (39.6-40.6)	94.5 (94.3-94.8)	83.6 (83.2-84.0)	72.1 (71.6-72.5)
75+	89.8 (89.5-90.1)	72.4 (72.0-72.8)	56.2 (55.7-56.6)	23.4 (23.0-23.8)	90.6 (90.4-90.9)	74.2 (73.8-74.6)	58.5 (58.0-58.9)
Men	94.8 (94.6-94.9)	84.2 (84.0-84.5)	73.0 (72.7-73.3)	43.7 (43.3-44.1)	95.2 (95.1-95.3)	85.5 (85.3-85.7)	74.9 (74.7-75.2)
Women	94.7 (94.5-94.8)	84.7 (84.4-85.0)	74.3 (73.9-74.7)	47.3 (46.8-47.8)	95.3 (95.1-95.4)	85.9 (85.7-86.2)	76.2 (75.8-76.5)
Diabetes	95.1 (94.9-95.3)	83.9 (83.6-84.3)	70.9 (70.4-71.4)	38.0 (37.4-38.6)	95.7 (95.5-95.9)	85.8 (85.5-86.2)	74.1 (73.7-74.6)
Hypertension / renal vascular disease	95.7 (95.5-95.9)	86.3 (85.9-86.6)	75.5 (75.0-75.9)	46.7 (46.1-47.3)	95.9 (95.7-96.1)	87.4 (87.1-87.7)	77.2 (76.8-77.6)
Glomerulonephritis	96.7 (96.4-97.0)	89.7 (89.2-90.2)	81.5 (80.8-82.2)	56.3 (55.3-57.4)	96.7 (96.4-97.0)	89.9 (89.4-90.4)	82.1 (81.5-82.8)
Other causes	93.5 (93.3-93.7)	82.3 (82.0-82.6)	71.8 (71.4-72.1)	45.4 (44.9-45.8)	94.2 (94.1-94.4)	83.6 (83.3-83.8)	73.4 (73.1-73.8)
All	94.7 (94.6-94.8)	84.4 (84.2-84.6)	73.5 (73.3-73.7)	45.0 (44.7-45.4)	95.2 (95.1-95.3)	85.7 (85.5-85.9)	75.4 (75.2-75.6)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)						
	cohort 2006 - 2010				cohort 2009 - 2013		
	90 day	1 year	2 year	5 year	90 day	1 year	2 year
0-19							
20-44	98.6 (97.6-99.6)	95.1 (93.1-97.1)	91.3 (88.6-94.1)	82.0 (77.6-86.7)	99.1 (98.3-100.0)	95.0 (92.9-97.1)	91.9 (89.2-94.6)
45-64	95.0 (94.1-96.0)	87.4 (85.9-88.9)	78.7 (76.8-80.6)	53.1 (50.5-55.8)	96.1 (95.2-97.0)	89.1 (87.7-90.6)	81.3 (79.5-83.2)
65-74	92.3 (91.0-93.5)	77.1 (75.2-79.1)	62.2 (59.9-64.5)	31.6 (29.3-34.0)	93.7 (92.6-94.8)	79.5 (77.6-81.4)	65.1 (62.8-67.4)
75+	87.7 (86.2-89.3)	67.0 (64.7-69.3)	47.4 (45.0-49.9)	15.8 (14.1-17.7)	90.6 (89.2-92.0)	71.6 (69.4-73.8)	53.7 (51.3-56.2)
Men	93.3 (92.5-94.1)	80.4 (79.2-81.7)	66.4 (64.9-68.0)	35.0 (33.2-36.9)	94.7 (94.0-95.4)	83.2 (82.0-84.4)	70.8 (69.4-72.3)
Women	92.8 (91.8-93.8)	80.6 (79.0-82.2)	68.2 (66.3-70.1)	39.7 (37.5-42.1)	94.6 (93.6-95.5)	82.9 (81.4-84.5)	71.4 (69.5-73.4)
Diabetes	93.8 (92.8-94.8)	79.7 (78.0-81.5)	63.0 (60.9-65.3)	30.9 (28.6-33.2)	95.7 (94.7-96.6)	83.0 (81.3-84.8)	68.4 (66.2-70.7)
Hypertension / renal vascular disease	94.6 (93.5-95.6)	82.6 (80.7-84.4)	68.8 (66.5-71.2)	38.7 (35.9-41.6)	95.7 (94.8-96.6)	85.2 (83.6-86.9)	73.6 (71.5-75.8)
Glomerulonephritis	95.9 (94.2-97.6)	88.6 (85.8-91.5)	78.6 (74.9-82.5)	46.6 (41.4-52.4)	95.9 (94.2-97.8)	90.7 (88.0-93.4)	82.3 (78.7-86.0)
Other causes	90.0 (88.7-91.3)	76.0 (74.1-77.9)	64.9 (62.7-67.1)	36.5 (34.0-39.2)	92.5 (91.4-93.6)	79.4 (77.7-81.1)	68.1 (66.0-70.2)
All	93.1 (92.5-93.7)	80.5 (79.5-81.5)	67.1 (65.9-68.3)	36.8 (35.4-38.2)	94.7 (94.1-95.2)	83.1 (82.2-84.1)	71.0 (69.9-72.2)

Table B.6.7
Incident dialysis patients, from day 91, unadjusted
by age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	96.6 (95.2-97.6)	94.1 (92.0-95.6)	89.3 (85.8-92.0)	97.0 (95.6-98.0)	94.5 (92.3-96.0)
20-44	96.5 (96.2-96.8)	92.8 (92.3-93.2)	81.8 (81.1-82.5)	96.7 (96.4-97.0)	93.3 (92.8-93.7)
45-64	90.6 (90.3-90.9)	82.5 (82.2-82.8)	58.8 (58.5-59.2)	91.3 (91.0-91.5)	83.6 (83.3-83.9)
65-74	83.3 (82.9-83.6)	70.5 (70.2-70.9)	39.7 (39.5-39.9)	85.0 (84.7-85.3)	72.6 (72.2-72.9)
75+	75.6 (75.3-75.9)	58.0 (57.8-58.3)	23.7 (23.6-23.8)	77.2 (76.9-77.5)	60.0 (59.8-60.3)
Men	84.3 (84.0-84.5)	71.7 (71.5-72.0)	41.2 (41.1-41.4)	85.3 (85.1-85.5)	72.9 (72.7-73.1)
Women	84.8 (84.5-85.0)	72.7 (72.4-73.0)	43.8 (43.6-44.0)	85.6 (85.3-85.9)	73.9 (73.6-74.1)
Diabetes	84.0 (83.7-84.4)	70.2 (69.8-70.6)	36.8 (36.6-37.0)	85.6 (85.3-85.9)	72.5 (72.1-72.9)
Hypertension / renal vascular disease	82.7 (82.3-83.1)	68.5 (68.1-68.9)	36.0 (35.7-36.2)	84.0 (83.7-84.4)	70.0 (69.7-70.4)
Glomerulonephritis	92.0 (91.6-92.4)	84.3 (83.7-84.8)	60.9 (60.3-61.5)	92.2 (91.8-92.6)	84.5 (83.9-85.0)
Other causes	84.0 (83.8-84.3)	72.5 (72.2-72.8)	44.8 (44.6-45.1)	84.6 (84.3-84.8)	72.9 (72.6-73.2)
All	84.4 (84.3-84.6)	72.1 (71.9-72.3)	42.2 (42.1-42.3)	85.4 (85.2-85.5)	73.2 (73.1-73.4)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	95.3 (93.0-96.8)	92.8 (90.1-94.8)	84.4 (80.5-87.6)	94.8 (92.3-96.5)	92.7 (89.9-94.7)
45-64	89.6 (88.2-90.8)	80.9 (79.3-82.4)	53.6 (52.2-55.1)	90.7 (89.3-91.9)	82.7 (81.1-84.3)
65-74	79.1 (77.4-80.6)	62.9 (61.3-64.4)	31.4 (30.7-32.2)	80.6 (78.9-82.1)	65.7 (64.1-67.3)
75+	69.8 (68.1-71.4)	49.4 (48.1-50.7)	15.7 (15.4-16.0)	73.5 (71.7-75.1)	54.9 (53.4-56.3)
Men	81.7 (80.6-82.8)	67.7 (66.6-68.8)	36.7 (36.0-37.3)	83.5 (82.4-84.6)	70.8 (69.6-71.9)
Women	81.2 (79.8-82.6)	67.5 (66.0-68.9)	38.0 (37.1-39.0)	82.3 (80.8-83.7)	69.2 (67.6-70.7)
Diabetes	79.5 (77.9-81.0)	62.8 (61.3-64.2)	29.6 (28.9-30.3)	81.0 (79.3-82.5)	66.3 (64.7-67.9)
Hypertension / renal vascular disease	79.7 (77.8-81.5)	63.4 (61.6-65.2)	32.5 (31.5-33.4)	82.2 (80.4-83.9)	68.1 (66.3-69.9)
Glomerulonephritis	91.7 (89.4-93.6)	84.6 (81.8-87.0)	59.2 (56.3-61.9)	93.3 (90.9-95.0)	86.5 (83.6-88.9)
Other causes	81.4 (79.8-82.9)	70.0 (68.4-71.6)	42.4 (41.2-43.5)	82.5 (81.0-83.9)	70.5 (68.9-72.0)
All	81.6 (80.7-82.4)	67.6 (66.8-68.5)	37.2 (36.6-37.7)	83.1 (82.2-83.9)	70.2 (69.3-71.1)

Table B.6.8
Incident dialysis patients, from day 91, adjusted
adjusted for age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	96.5 (95.2-97.8)	93.5 (91.3-95.7)	86.4 (81.6-91.4)	96.9 (95.6-98.2)	94.3 (92.3-96.3)
20-44	96.5 (96.1-96.8)	92.6 (92.1-93.1)	80.5 (79.5-81.4)	96.6 (96.3-97.0)	93.2 (92.7-93.6)
45-64	91.0 (90.7-91.3)	83.0 (82.6-83.4)	58.7 (58.1-59.3)	91.5 (91.2-91.8)	84.0 (83.6-84.4)
65-74	83.9 (83.5-84.3)	71.5 (71.0-72.0)	40.8 (40.2-41.3)	85.4 (85.1-85.8)	73.5 (73.0-73.9)
75+	76.0 (75.5-76.4)	58.5 (58.1-59.0)	24.2 (23.8-24.7)	77.5 (77.1-77.9)	60.7 (60.3-61.2)
Men	85.8 (85.6-86.1)	74.2 (73.9-74.5)	43.9 (43.5-44.3)	87.0 (86.8-87.2)	76.0 (75.7-76.3)
Women	86.7 (86.4-87.0)	75.8 (75.4-76.2)	47.9 (47.4-48.4)	87.6 (87.3-87.9)	77.4 (77.0-77.7)
Diabetes	84.7 (84.3-85.1)	71.2 (70.7-71.7)	37.6 (37.0-38.2)	86.5 (86.2-86.9)	74.3 (73.8-74.8)
Hypertension / renal vascular disease	87.2 (86.8-87.6)	75.9 (75.4-76.4)	46.5 (45.8-47.1)	88.5 (88.1-88.8)	77.8 (77.4-78.3)
Glomerulonephritis	90.7 (90.2-91.3)	82.0 (81.3-82.7)	56.3 (55.2-57.3)	91.0 (90.5-91.5)	82.6 (81.9-83.3)
Other causes	85.0 (84.7-85.3)	74.0 (73.7-74.4)	46.4 (46.0-46.9)	85.9 (85.6-86.2)	75.3 (74.9-75.6)
All	86.2 (86.0-86.3)	74.8 (74.5-75.0)	45.4 (45.1-45.7)	87.3 (87.1-87.4)	76.5 (76.2-76.7)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Castile-La Mancha), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	94.9 (92.7-97.0)	92.2 (89.6-95.0)	83.1 (78.7-87.8)	94.4 (92.1-96.7)	92.3 (89.6-95.0)
45-64	89.9 (88.4-91.3)	81.4 (79.6-83.3)	53.9 (51.2-56.8)	90.8 (89.4-92.2)	83.1 (81.2-85.0)
65-74	79.8 (77.9-81.8)	64.2 (61.9-66.6)	32.7 (30.3-35.3)	81.0 (79.1-82.9)	66.6 (64.3-69.0)
75+	70.4 (68.1-72.8)	50.2 (47.6-52.9)	16.6 (14.7-18.7)	73.8 (71.5-76.1)	55.6 (53.0-58.2)
Men	82.5 (81.2-83.8)	68.5 (66.9-70.1)	35.8 (34.0-37.8)	84.5 (83.3-85.7)	72.4 (70.8-73.9)
Women	83.5 (81.9-85.1)	70.8 (68.8-72.8)	40.8 (38.4-43.3)	84.6 (83.0-86.2)	72.7 (70.7-74.7)
Diabetes	80.5 (78.7-82.4)	64.2 (61.9-66.5)	30.0 (27.7-32.5)	82.2 (80.4-84.1)	68.4 (66.1-70.8)
Hypertension / renal vascular disease	83.7 (81.8-85.6)	69.4 (67.0-71.9)	39.1 (36.3-42.2)	86.1 (84.4-87.8)	74.4 (72.2-76.7)
Glomerulonephritis	89.1 (86.2-92.1)	79.9 (76.1-83.9)	47.7 (42.4-53.7)	91.5 (88.8-94.2)	83.1 (79.4-87.0)
Other causes	81.2 (79.3-83.1)	69.2 (66.9-71.5)	38.7 (36.0-41.7)	82.9 (81.2-84.6)	70.9 (68.8-73.1)
All	82.9 (81.9-83.9)	69.3 (68.1-70.6)	37.7 (36.2-39.2)	84.5 (83.6-85.5)	72.5 (71.3-73.7)

Table B.6.9
First transplant patients (deceased donor), from day of transplant, unadjusted
by age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	99.2 (98.2-99.6)	99.0 (98.0-99.5)	97.6 (96.3-98.5)	98.4 (97.1-99.1)	97.9 (96.6-98.7)
20-44	98.4 (98.1-98.7)	97.7 (97.3-98.0)	95.3 (94.9-95.8)	98.7 (98.4-98.9)	98.1 (97.7-98.3)
45-64	96.5 (96.2-96.8)	94.6 (94.2-95.0)	88.1 (87.6-88.6)	96.8 (96.5-97.0)	94.8 (94.5-95.1)
65+	91.0 (90.2-91.7)	87.4 (86.5-88.2)	74.1 (73.1-75.0)	92.5 (91.9-93.0)	88.5 (87.8-89.1)
Men	95.9 (95.6-96.2)	93.9 (93.5-94.2)	86.9 (86.4-87.3)	96.2 (95.9-96.4)	93.9 (93.6-94.2)
Women	96.6 (96.2-96.9)	95.1 (94.7-95.5)	89.7 (89.2-90.3)	96.5 (96.2-96.8)	94.9 (94.6-95.3)
Diabetes	94.6 (93.9-95.2)	92.1 (91.3-92.8)	83.1 (82.1-84.0)	94.6 (94.0-95.1)	91.6 (90.8-92.2)
Hypertension / renal vascular disease	93.7 (92.7-94.5)	90.8 (89.8-91.8)	81.8 (80.6-83.0)	94.7 (94.0-95.3)	91.7 (90.9-92.5)
Glomerulonephritis	97.2 (96.7-97.6)	96.0 (95.5-96.5)	91.1 (90.4-91.8)	97.6 (97.2-98.0)	96.2 (95.7-96.6)
Other causes	96.7 (96.4-97.0)	95.1 (94.7-95.4)	89.5 (89.0-89.9)	96.8 (96.5-97.0)	95.0 (94.7-95.3)
All	96.1 (95.9-96.4)	94.3 (94.1-94.6)	88.0 (87.6-88.3)	96.3 (96.1-96.5)	94.3 (94.0-94.5)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	96.6 (94.0-98.1)	95.6 (92.8-97.4)	92.6 (89.3-94.9)	98.1 (95.6-99.2)	97.4 (94.7-98.7)
45-64	94.6 (92.6-96.0)	92.1 (90.0-93.9)	86.0 (83.5-88.2)	95.9 (94.1-97.1)	94.2 (92.3-95.7)
65+	86.0 (82.2-89.1)	79.9 (75.9-83.3)	67.9 (64.1-71.4)	90.6 (87.5-93.0)	85.9 (82.5-88.7)
Men	92.9 (91.1-94.4)	89.4 (87.4-91.2)	82.1 (79.8-84.1)	94.3 (92.7-95.6)	91.4 (89.5-93.0)
Women	93.7 (91.2-95.6)	91.9 (89.1-93.9)	86.5 (83.4-89.1)	96.1 (94.0-97.5)	95.0 (92.7-96.6)
Diabetes	88.5 (84.6-91.4)	85.5 (81.5-88.7)	78.8 (74.7-82.4)	91.0 (87.4-93.6)	87.8 (84.0-90.8)
Hypertension / renal vascular disease	91.0 (85.9-94.3)	85.8 (80.3-89.9)	72.9 (67.4-77.6)	95.7 (91.8-97.8)	93.0 (88.7-95.7)
Glomerulonephritis	95.7 (92.9-97.4)	92.4 (89.1-94.7)	87.1 (83.4-90.1)	97.0 (94.5-98.4)	94.7 (91.7-96.6)
Other causes	94.8 (92.6-96.3)	92.9 (90.5-94.7)	87.2 (84.5-89.5)	95.6 (93.6-97.0)	93.9 (91.7-95.5)
All	93.2 (91.7-94.4)	90.3 (88.7-91.7)	83.6 (81.8-85.2)	95.0 (93.7-96.0)	92.7 (91.2-93.9)

Table B.6.10
First transplant patients (deceased donor), from day of transplant, adjusted
adjusted for age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	99.1 (98.4-99.8)	98.9 (98.2-99.7)	97.4 (96.2-98.6)	98.5 (97.6-99.4)	98.0 (97.0-99.1)
20-44	98.5 (98.2-98.7)	97.8 (97.4-98.1)	95.6 (95.1-96.1)	98.8 (98.5-99.0)	98.1 (97.8-98.4)
45-64	96.7 (96.4-97.0)	94.9 (94.5-95.2)	88.6 (88.1-89.2)	96.9 (96.7-97.2)	95.1 (94.8-95.4)
65+	91.6 (90.9-92.4)	88.2 (87.2-89.1)	75.5 (74.3-76.8)	93.2 (92.6-93.7)	89.4 (88.7-90.1)
Men	97.3 (97.1-97.6)	96.0 (95.7-96.3)	91.1 (90.7-91.6)	97.8 (97.6-98.0)	96.4 (96.1-96.6)
Women	97.6 (97.4-97.9)	96.6 (96.2-96.9)	92.6 (92.2-93.1)	97.8 (97.6-98.1)	96.8 (96.5-97.1)
Diabetes	96.1 (95.6-96.6)	94.3 (93.6-94.9)	87.3 (86.4-88.2)	96.6 (96.2-97.0)	94.6 (94.1-95.2)
Hypertension / renal vascular disease	96.7 (96.2-97.2)	95.2 (94.6-95.8)	90.1 (89.2-91.0)	97.4 (97.1-97.8)	95.9 (95.4-96.4)
Glomerulonephritis	98.0 (97.6-98.3)	97.1 (96.7-97.5)	93.4 (92.8-94.0)	98.4 (98.1-98.6)	97.4 (97.1-97.7)
Other causes	97.6 (97.4-97.9)	96.4 (96.1-96.7)	92.1 (91.7-92.5)	97.9 (97.7-98.1)	96.7 (96.4-96.9)
All	97.5 (97.3-97.6)	96.2 (96.0-96.4)	91.7 (91.3-92.1)	97.8 (97.7-98.0)	96.5 (96.3-96.7)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	96.9 (95.0-98.8)	95.9 (93.8-98.1)	93.1 (90.4-95.9)	98.2 (96.7-99.8)	97.5 (95.7-99.3)
45-64	95.0 (93.4-96.6)	92.8 (90.8-94.7)	87.1 (84.6-89.7)	96.2 (94.8-97.6)	94.6 (93.0-96.3)
65+	87.5 (84.0-91.2)	81.6 (77.5-85.9)	70.8 (66.0-76.0)	91.8 (89.2-94.5)	87.4 (84.2-90.6)
Men	95.9 (94.8-96.9)	93.7 (92.4-95.0)	89.0 (87.3-90.8)	97.1 (96.2-97.9)	95.4 (94.4-96.5)
Women	96.0 (94.6-97.5)	94.8 (93.2-96.5)	91.2 (89.1-93.4)	97.7 (96.6-98.7)	97.0 (95.8-98.2)
Diabetes	92.7 (90.2-95.2)	90.7 (87.9-93.5)	86.0 (82.6-89.5)	95.1 (93.2-97.0)	93.2 (91.0-95.4)
Hypertension / renal vascular disease	95.8 (93.6-98.0)	92.9 (90.1-95.7)	86.3 (82.5-90.3)	97.9 (96.5-99.4)	96.3 (94.5-98.2)
Glomerulonephritis	97.0 (95.4-98.6)	94.6 (92.5-96.8)	90.6 (87.8-93.5)	98.1 (96.9-99.3)	96.6 (95.0-98.3)
Other causes	96.5 (95.1-97.8)	95.1 (93.6-96.7)	91.0 (88.9-93.1)	97.2 (96.1-98.3)	96.1 (94.8-97.4)
All	96.0 (95.1-96.8)	94.1 (93.1-95.2)	89.8 (88.5-91.2)	97.3 (96.6-97.9)	96.0 (95.2-96.8)

Table B.6.11
First transplant patients (living donor), from day of transplant, unadjusted
by age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	99.6 (98.2-99.9)	99.6 (98.2-99.9)	99.3 (97.9-99.8)	99.8 (98.5-100.0)	99.8 (98.5-100.0)
20-44	99.4 (99.1-99.6)	99.2 (98.8-99.4)	97.6 (97.1-98.1)	99.7 (99.5-99.9)	99.4 (99.1-99.6)
45-64	98.1 (97.6-98.5)	97.0 (96.4-97.5)	92.6 (91.8-93.4)	98.5 (98.1-98.8)	97.4 (97.0-97.8)
65+	95.6 (93.9-96.8)	92.4 (90.4-94.1)	80.0 (77.5-82.3)	96.9 (95.7-97.8)	93.6 (92.0-94.8)
Men	98.5 (98.1-98.8)	97.7 (97.2-98.0)	93.9 (93.3-94.5)	98.8 (98.6-99.1)	97.9 (97.6-98.3)
Women	98.7 (98.2-99.0)	97.7 (97.2-98.2)	94.4 (93.6-95.1)	99.0 (98.6-99.3)	97.9 (97.4-98.3)
Diabetes	96.4 (94.7-97.5)	94.0 (92.1-95.5)	85.4 (82.9-87.5)	97.2 (95.9-98.1)	95.5 (94.0-96.7)
Hypertension / renal vascular disease	96.9 (95.4-98.0)	95.9 (94.1-97.1)	89.6 (87.3-91.5)	97.9 (96.7-98.6)	96.4 (94.9-97.4)
Glomerulonephritis	99.2 (98.7-99.5)	98.7 (98.2-99.1)	96.6 (95.8-97.3)	99.3 (98.9-99.5)	98.6 (98.1-99.0)
Other causes	98.9 (98.6-99.2)	98.2 (97.8-98.5)	95.0 (94.4-95.6)	99.1 (98.9-99.3)	98.2 (97.9-98.5)
All	98.6 (98.3-98.8)	97.7 (97.4-98.0)	94.1 (93.6-94.6)	98.9 (98.7-99.1)	97.9 (97.6-98.2)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	100	100	99.0 (93.5-99.9)	100	100
45-64	97.1 (91.5-99.0)	95.1 (89.1-97.8)	90.1 (83.4-94.2)	97.3 (92.2-99.1)	96.4 (91.0-98.6)
65+					
Men	98.8 (95.5-99.7)	98.3 (94.8-99.4)	94.2 (89.8-96.7)	98.4 (95.2-99.5)	97.9 (94.6-99.2)
Women					
Diabetes					
Hypertension / renal vascular disease					
Glomerulonephritis					
Other causes	98.6 (94.6-99.6)	97.9 (93.7-99.3)	93.6 (88.6-96.5)	99.3 (95.4-99.9)	99.3 (95.4-99.9)
All	98.8 (96.5-99.6)	98.1 (95.5-99.2)	94.5 (91.2-96.6)	98.9 (96.7-99.6)	98.5 (96.2-99.4)

Table B.6.12
First transplant patients (living donor), from day of transplant, adjusted
adjusted for age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	99.5 (98.7-100.0)	99.5 (98.7-100.0)	99.2 (98.3-100.0)	99.8 (99.3-100.0)	99.8 (99.3-100.0)
20-44	99.4 (99.1-99.7)	99.1 (98.8-99.4)	97.5 (97.0-98.0)	99.7 (99.6-99.9)	99.3 (99.1-99.6)
45-64	98.3 (97.9-98.7)	97.2 (96.7-97.7)	92.9 (92.0-93.7)	98.6 (98.3-98.9)	97.5 (97.1-98.0)
65+	96.2 (94.8-97.6)	93.6 (91.8-95.4)	82.1 (79.3-85.0)	96.9 (95.8-97.9)	93.8 (92.4-95.3)
Men	98.8 (98.5-99.0)	98.1 (97.7-98.4)	94.6 (93.9-95.2)	99.1 (98.8-99.3)	98.4 (98.0-98.7)
Women	98.7 (98.3-99.1)	97.8 (97.2-98.3)	94.4 (93.5-95.2)	99.1 (98.8-99.4)	98.0 (97.6-98.5)
Diabetes	97.3 (96.2-98.3)	95.5 (94.1-96.9)	88.6 (86.5-90.8)	98.1 (97.3-98.9)	96.9 (95.9-97.9)
Hypertension / renal vascular disease	98.1 (97.2-98.9)	97.4 (96.4-98.4)	93.1 (91.5-94.8)	98.7 (98.1-99.3)	97.8 (97.0-98.6)
Glomerulonephritis	99.1 (98.7-99.5)	98.6 (98.1-99.1)	96.3 (95.4-97.1)	99.2 (98.9-99.6)	98.5 (98.1-99.0)
Other causes	98.9 (98.6-99.2)	98.2 (97.8-98.6)	94.9 (94.2-95.6)	99.2 (99.0-99.4)	98.4 (98.0-98.7)
All	98.7 (98.5-99.0)	98.0 (97.7-98.3)	94.5 (94.0-95.0)	99.1 (98.9-99.2)	98.2 (98.0-98.5)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	100	100	98.9 (96.7-100.0)		
45-64	97.2 (94.1-100.0)	95.3 (91.3-99.4)	90.5 (85.0-96.3)	97.1 (94.0-100.0)	96.2 (92.6-99.9)
65+					
Men	98.9 (97.3-100.0)	98.3 (96.4-100.0)	94.1 (90.6-97.7)	98.6 (97.0-100.0)	98.1 (96.3-100.0)
Women					
Diabetes					
Hypertension / renal vascular disease					
Glomerulonephritis					
Other causes	98.4 (96.3-100.0)	97.6 (95.0-100.0)	92.7 (88.3-97.4)	99.3 (97.8-100.0)	99.3 (97.8-100.0)
All	98.8 (97.5-100.0)	98.0 (96.3-99.8)	94.2 (91.4-97.2)	98.9 (97.7-100.0)	98.5 (97.1-100.0)

Table B.6.13
First graft (deceased donor), from day of transplant, unadjusted
by age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	92.7 (90.7-94.3)	90.0 (87.9-91.8)	82.9 (80.5-85.0)	91.8 (89.7-93.4)	89.3 (87.1-91.1)
20-44	93.8 (93.2-94.3)	91.8 (91.2-92.4)	84.8 (84.1-85.5)	94.3 (93.8-94.8)	92.3 (91.8-92.9)
45-64	91.3 (90.8-91.7)	88.6 (88.1-89.1)	79.9 (79.4-80.4)	91.7 (91.3-92.1)	88.9 (88.5-89.3)
65+	84.7 (83.8-85.6)	80.6 (79.6-81.5)	65.9 (65.0-66.8)	86.0 (85.3-86.6)	81.3 (80.5-82.0)
Men	90.9 (90.5-91.3)	88.1 (87.6-88.5)	78.5 (78.0-79.0)	91.0 (90.6-91.3)	87.7 (87.3-88.1)
Women	90.8 (90.2-91.3)	88.3 (87.7-88.8)	79.7 (79.0-80.3)	91.2 (90.7-91.7)	88.7 (88.2-89.2)
Diabetes	90.2 (89.3-91.0)	86.9 (85.9-87.7)	75.6 (74.6-76.6)	89.7 (89.0-90.4)	85.8 (85.0-86.6)
Hypertension / renal vascular disease	88.2 (87.1-89.2)	84.9 (83.7-86.0)	72.1 (70.8-73.3)	88.7 (87.7-89.5)	84.9 (83.9-85.8)
Glomerulonephritis	91.3 (90.6-92.0)	88.9 (88.2-89.6)	80.7 (79.8-81.5)	91.9 (91.3-92.5)	89.2 (88.6-89.9)
Other causes	91.4 (91.0-91.9)	88.9 (88.4-89.4)	80.6 (80.1-81.2)	91.7 (91.3-92.1)	89.1 (88.6-89.5)
All	90.9 (90.5-91.2)	88.1 (87.8-88.5)	78.9 (78.5-79.3)	91.1 (90.8-91.3)	88.1 (87.7-88.4)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	92.3 (88.9-94.6)	89.6 (86.0-92.3)	79.1 (75.2-82.5)	92.9 (89.4-95.3)	88.4 (84.5-91.4)
45-64	89.4 (87.1-91.4)	85.3 (82.7-87.5)	79.0 (76.3-81.4)	91.8 (89.6-93.5)	88.5 (86.2-90.5)
65+	81.6 (77.6-84.9)	76.1 (72.1-79.6)	62.8 (59.2-66.2)	84.5 (81.1-87.4)	79.0 (75.4-82.1)
Men	88.8 (86.8-90.6)	84.5 (82.3-86.5)	75.6 (73.2-77.7)	89.6 (87.6-91.3)	85.4 (83.2-87.3)
Women	87.7 (84.7-90.1)	84.2 (81.0-86.9)	74.9 (71.7-77.8)	90.9 (88.2-93.0)	87.0 (84.0-89.4)
Diabetes	83.3 (79.2-86.7)	78.8 (74.7-82.4)	71.4 (67.3-75.1)	87.8 (84.0-90.8)	82.1 (78.0-85.4)
Hypertension / renal vascular disease	88.4 (83.0-92.1)	82.6 (77.0-86.9)	69.0 (63.7-73.7)	89.8 (85.1-93.1)	83.3 (78.3-87.3)
Glomerulonephritis	90.1 (86.6-92.7)	85.5 (81.7-88.5)	75.9 (72.0-79.3)	90.1 (86.6-92.7)	86.4 (82.7-89.4)
Other causes	90.2 (87.6-92.2)	87.3 (84.5-89.5)	78.9 (76.0-81.5)	91.3 (88.9-93.3)	88.6 (86.0-90.7)
All	88.4 (86.8-89.9)	84.4 (82.6-86.0)	75.3 (73.5-77.1)	90.1 (88.5-91.4)	85.9 (84.2-87.5)

Table B.6.14
First graft (deceased donor), from day of transplant, adjusted
adjusted for age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	92.2 (90.2-94.2)	89.4 (87.1-91.7)	81.7 (78.8-84.7)	91.5 (89.4-93.6)	88.8 (86.4-91.2)
20-44	93.8 (93.3-94.4)	91.9 (91.3-92.6)	85.1 (84.3-85.9)	94.3 (93.8-94.9)	92.4 (91.8-93.0)
45-64	91.4 (90.9-91.9)	88.8 (88.3-89.3)	80.2 (79.5-80.8)	91.8 (91.4-92.3)	89.1 (88.6-89.6)
65+	85.3 (84.3-86.3)	81.2 (80.0-82.3)	66.9 (65.5-68.3)	86.6 (85.8-87.4)	82.0 (81.1-82.9)
Men	91.9 (91.5-92.3)	89.4 (88.9-89.8)	80.5 (79.9-81.2)	92.4 (92.1-92.8)	89.6 (89.2-90.1)
Women	91.5 (90.9-92.0)	89.1 (88.5-89.7)	81.0 (80.2-81.8)	92.3 (91.8-92.8)	90.0 (89.5-90.5)
Diabetes	90.9 (90.0-91.7)	87.8 (86.8-88.8)	77.0 (75.7-78.3)	91.2 (90.5-91.9)	87.7 (86.9-88.6)
Hypertension / renal vascular disease	90.6 (89.6-91.6)	87.8 (86.6-88.9)	77.0 (75.5-78.5)	91.4 (90.6-92.2)	88.3 (87.4-89.3)
Glomerulonephritis	91.8 (91.2-92.6)	89.5 (88.8-90.3)	81.7 (80.7-82.7)	92.8 (92.2-93.4)	90.3 (89.6-91.0)
Other causes	92.0 (91.5-92.5)	89.6 (89.1-90.1)	81.7 (81.0-82.4)	92.6 (92.2-93.0)	90.2 (89.8-90.7)
All	91.8 (91.4-92.1)	89.3 (88.9-89.7)	80.7 (80.2-81.2)	92.4 (92.1-92.7)	89.8 (89.4-90.1)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	92.6 (89.7-95.5)	90.0 (86.7-93.4)	79.8 (75.5-84.5)	93.0 (90.1-96.1)	88.6 (84.9-92.5)
45-64	89.9 (87.6-92.2)	85.8 (83.2-88.5)	79.7 (76.7-82.9)	91.9 (89.9-94.0)	89.0 (86.7-91.4)
65+	82.7 (78.6-87.0)	77.2 (72.7-82.0)	64.7 (59.6-70.3)	85.7 (82.3-89.2)	80.2 (76.3-84.3)
Men	90.7 (88.9-92.6)	86.9 (84.8-89.1)	79.2 (76.6-81.8)	91.8 (90.1-93.4)	88.3 (86.4-90.3)
Women	89.1 (86.4-91.9)	86.0 (82.9-89.1)	77.4 (73.7-81.3)	92.2 (90.0-94.5)	88.8 (86.2-91.5)
Diabetes	85.4 (81.5-89.4)	81.4 (77.1-85.9)	74.6 (69.8-79.7)	90.2 (87.1-93.4)	85.4 (81.7-89.2)
Hypertension / renal vascular disease	91.0 (87.2-95.1)	85.9 (81.2-90.9)	75.3 (69.4-81.6)	92.2 (89.0-95.7)	86.8 (82.7-91.2)
Glomerulonephritis	90.9 (87.9-94.1)	86.6 (83.0-90.4)	77.5 (73.1-82.2)	91.4 (88.5-94.4)	88.2 (84.9-91.7)
Other causes	91.1 (88.7-93.4)	88.4 (85.8-91.1)	80.6 (77.4-83.9)	92.2 (90.1-94.3)	89.9 (87.5-92.3)
All	90.2 (88.7-91.7)	86.6 (84.9-88.4)	78.6 (76.5-80.8)	92.0 (90.6-93.3)	88.5 (87.0-90.1)

Table B.6.15
First graft (living donor), from day of transplant, unadjusted
by age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	97.1 (95.1-98.3)	96.2 (94.1-97.6)	89.4 (86.6-91.7)	96.8 (94.8-98.0)	96.4 (94.3-97.7)
20-44	96.4 (95.8-96.9)	94.7 (94.0-95.4)	89.0 (88.0-89.8)	97.3 (96.8-97.8)	95.8 (95.2-96.4)
45-64	95.8 (95.2-96.4)	94.1 (93.3-94.8)	87.8 (86.8-88.7)	96.1 (95.5-96.6)	94.2 (93.5-94.8)
65+	92.6 (90.6-94.2)	89.2 (86.9-91.0)	75.3 (72.8-77.6)	95.2 (93.8-96.3)	91.5 (89.8-92.9)
Men	95.9 (95.3-96.3)	94.3 (93.7-94.9)	87.5 (86.6-88.2)	96.6 (96.1-97.0)	95.0 (94.5-95.5)
Women	95.9 (95.2-96.5)	93.7 (92.8-94.4)	87.2 (86.1-88.2)	96.4 (95.8-96.9)	94.1 (93.3-94.7)
Diabetes	93.6 (91.6-95.1)	90.7 (88.5-92.5)	79.3 (76.7-81.5)	94.9 (93.3-96.2)	92.4 (90.6-93.9)
Hypertension / renal vascular disease	93.9 (91.9-95.4)	91.4 (89.3-93.2)	82.1 (79.5-84.3)	95.4 (93.9-96.6)	93.2 (91.4-94.6)
Glomerulonephritis	95.9 (95.0-96.6)	94.2 (93.2-95.0)	87.8 (86.5-89.0)	96.5 (95.7-97.1)	94.7 (93.9-95.5)
Other causes	96.5 (96.0-97.0)	95.0 (94.3-95.5)	89.2 (88.3-89.9)	96.8 (96.4-97.3)	95.1 (94.6-95.6)
All	95.9 (95.4-96.3)	94.1 (93.6-94.5)	87.4 (86.7-88.0)	96.5 (96.2-96.8)	94.7 (94.3-95.1)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	98.1 (92.8-99.5)	96.2 (90.4-98.5)	91.3 (84.8-95.1)	96.4 (90.9-98.6)	95.5 (89.8-98.0)
45-64	96.1 (90.3-98.5)	93.1 (86.8-96.5)	84.2 (77.1-89.3)	95.5 (90.0-98.0)	94.6 (88.9-97.4)
65+					
Men	97.1 (93.4-98.8)	96.0 (91.9-98.0)	88.4 (83.3-92.0)	96.3 (92.6-98.2)	95.2 (91.3-97.4)
Women					
Diabetes					
Hypertension / renal vascular disease					
Glomerulonephritis					
Other causes	98.6 (94.6-99.6)	97.2 (92.9-98.9)	90.1 (84.6-93.7)	98.0 (94.0-99.3)	97.3 (93.2-99.0)
All	97.3 (94.5-98.7)	95.3 (92.2-97.3)	88.3 (84.3-91.3)	96.3 (93.4-98.0)	95.6 (92.6-97.4)

Table B.6.16
First graft (living donor), from day of transplant, adjusted
adjusted for age, gender, and primary renal disease

All Countries *	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19	96.6 (94.8-98.4)	95.8 (93.8-97.8)	88.1 (84.9-91.4)	96.4 (94.6-98.2)	95.9 (94.0-97.8)
20-44	96.3 (95.6-96.9)	94.6 (93.8-95.3)	88.5 (87.5-89.6)	97.2 (96.7-97.7)	95.7 (95.1-96.3)
45-64	95.9 (95.2-96.5)	94.1 (93.4-94.9)	87.8 (86.7-88.9)	96.1 (95.5-96.6)	94.2 (93.5-94.9)
65+	92.9 (91.0-94.8)	89.8 (87.6-92.1)	76.4 (73.3-79.8)	94.9 (93.5-96.3)	91.2 (89.5-93.0)
Men	95.8 (95.2-96.4)	94.2 (93.6-94.9)	87.0 (86.1-88.0)	96.4 (96.0-96.9)	94.9 (94.4-95.5)
Women	95.4 (94.6-96.1)	92.9 (92.0-93.9)	85.8 (84.5-87.1)	96.0 (95.3-96.6)	93.4 (92.6-94.3)
Diabetes	93.7 (92.0-95.5)	90.9 (88.7-93.0)	79.6 (76.6-82.6)	95.2 (93.7-96.6)	92.7 (91.0-94.5)
Hypertension / renal vascular disease	94.1 (92.4-95.9)	91.8 (89.7-93.9)	82.7 (79.9-85.6)	95.7 (94.4-97.0)	93.6 (92.0-95.2)
Glomerulonephritis	95.5 (94.6-96.4)	93.7 (92.6-94.8)	86.8 (85.3-88.4)	96.2 (95.4-96.9)	94.3 (93.4-95.2)
Other causes	96.3 (95.7-96.8)	94.7 (94.0-95.3)	88.5 (87.5-89.4)	96.7 (96.2-97.2)	94.9 (94.4-95.5)
All	95.6 (95.2-96.1)	93.8 (93.2-94.3)	86.6 (85.8-87.4)	96.3 (95.9-96.7)	94.4 (93.9-94.9)

* Based on data from Austria, Belgium (Dutch-speaking), Belgium (French-speaking), Denmark, Finland, France, Greece, Iceland, Norway, Spain (Andalusia), Spain (Aragon), Spain (Asturias), Spain (Basque country), Spain (Cantabria), Spain (Castile and León), Spain (Catalonia), Spain (Extremadura), Spain (Galicia), Spain (Community of Madrid), Spain (Valencian region), Sweden, the Netherlands, United Kingdom (England/Northern Ireland/Wales) and United Kingdom (Scotland)

Austria	Survival probabilities as % (95% CI)				
	cohort 2006 - 2010			cohort 2009 - 2013	
	1 year	2 year	5 year	1 year	2 year
0-19					
20-44	98.0 (95.2-100.0)	95.9 (92.0-99.9)	90.7 (85.1-96.7)	96.2 (92.7-99.9)	95.3 (91.3-99.4)
45-64	96.2 (92.7-99.9)	93.4 (88.7-98.2)	84.7 (78.1-91.9)	95.4 (91.5-99.4)	94.4 (90.2-98.9)
65+					
Men	96.9 (94.3-99.6)	95.6 (92.5-98.9)	87.4 (82.4-92.8)	96.1 (93.3-99.0)	95.0 (91.8-98.2)
Women					
Diabetes					
Hypertension / renal vascular disease					
Glomerulonephritis					
Other causes	98.4 (96.3-100.0)	96.9 (93.9-99.9)	89.0 (83.7-94.6)	97.8 (95.3-100.0)	97.0 (94.2-99.9)
All	97.0 (94.8-99.2)	94.8 (92.0-97.7)	87.0 (82.7-91.4)	95.9 (93.5-98.4)	95.1 (92.4-97.8)